

Diabetics do not benefit from implantable defibrillator pacemakers

November 21 2019



Credit: CC0 Public Domain

According to a broad European study involving multiple medical institutes, patients with diabetes do not benefit from implantable cardioverter defibrillator (ICD) pacemakers.

Current treatment recommendations support ICD pacemaker treatment for all patients with severe malfunction of the left ventricle. The purpose of the defibrillator is to protect the patient from life-threatening cardiac arrhythmias. In practice, the device detects a [cardiac arrhythmia](#) and corrects it either by overdrive pacing or by applying a small shock.

In Finland, around 900 ICD pacemakers are implanted per year to prevent cardiac arrest.

However, the benefits of the device would appear to be small for diabetes patients. The results are surprising, as heart patients with diabetes are known to face a significantly higher risk of sudden death than other heart patients.

"Surprisingly, diabetes patients did not seem to benefit as much as patients who did not have diabetes," says Professor Juhani Junttila from the University of Oulu, Finland.

In diabetes patients, the prevention of death and incidence of arrhythmia corrections were significantly lower compared to patients without [diabetes](#).

Diabetes patients with severe dysfunction of the left ventricle have considerably more [health problems](#) than other patients with heart dysfunction, and so they already face a poor prognosis. According to Junttila, this is a probable reason why the ICD pacemaker does not bring much benefit.

The results of the study suggest that assessing dysfunction by measuring cardiac pumping capacity is not sufficient for assessing the benefits of implanting an ICD pacemaker.

"ICD pacemaker treatment is effective when applied to the correct

patient group, but with the current patient selections it does not, however, prevent the majority of sudden deaths. New tools should therefore be developed for patient selection so that ICD pacemaker treatment can be better allocated," Junttila says.

Data for a total of 3,535 patients with an ICD pacemakers was collected for this international EU-CERT study. The study was recently published in *Diabetes Care*.

There are currently extensive further studies being carried out in both Europe and the United States on the usefulness of ICD [pacemaker](#) treatment for [diabetes patients](#).

More information: M. Juhani Junttila et al. Appropriate Shocks and Mortality in Patients With Versus Without Diabetes With Prophylactic Implantable Cardioverter Defibrillators, *Diabetes Care* (2019). [DOI: 10.2337/dc19-1014](#)

Provided by University of Oulu

Citation: Diabetics do not benefit from implantable defibrillator pacemakers (2019, November 21) retrieved 25 April 2024 from <https://medicalxpress.com/news/2019-11-diabetics-benefit-implantable-defibrillator-pacemakers.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--