


# Researchers explore neuromarkers for poor social outcomes after traumatic brain injury

November 7 2019

---



Dr. Lengenfelder is assistant director of the Center for Traumatic Brain Injury Research at Kessler Foundation. Credit: Kessler Foundation/Jody Banks

A recent study of individuals with moderate to severe brain injury yielded new information linking brain connectivity patterns and deficits of emotion recognition and social functioning. The article, "Default mode network connectivity predicts emotion recognition and social integration after traumatic brain injury", was published on August 9,

2019 by *Frontiers in Neurology*.

A common consequence of moderate to severe [traumatic brain injury](#) (TBI) is a deficit in the ability to recognize the emotions of others. This, in turn, interferes with the ability to function effectively in [social settings](#) including school, work, and the community. Brain imaging studies have shown that emotion recognition ability may be linked to connectivity (i.e., coordinated activity) between regions in a functional network known as the default mode network. However, little research has been done on the relationship between resting-state functional connectivity and emotion recognition in individuals with or without TBI. This study is the first to explore these relationships and examine their potential value as predictors of social outcomes after TBI.

The researchers conducted tests of emotion recognition in two groups, 21 individuals with TBI and 27 healthy controls, and correlated their performance with resting-state functional connectivity between regions of the default mode network. They found that for all participants, deficits in emotion recognition were linked to increased connectivity between prefrontal and temporal regions of the network. In the group with TBI, the strength of prefrontal-temporal connectivity predicted participants' level of social integration using the Community Integration Questionnaire, an important measure of [social functioning](#) post-TBI.

"This study advances our understanding of the brain basis of post-TBI deficits in social functioning," said Dr. Lancaster, the Hearst Postdoctoral Fellow in the Center for Traumatic Brain Injury Research at Kessler Foundation. "In particular, we found that default mode network connectivity was a stronger predictor of social outcomes than [emotion recognition](#) alone," she noted, "underscoring the importance of brain connectivity metrics in explaining social aspects of TBI recovery."

Further research is needed to confirm the reliability and validity of

increased connectivity as a neuromarker for post-TBI social functioning. Neuromarkers will help clinicians identify individuals at risk for poor outcomes after TBI, and encourage the development of new rehabilitative interventions aimed at improving [social integration](#) and quality of life.

**More information:** Katie Lancaster et al, Default Mode Network Connectivity Predicts Emotion Recognition and Social Integration After Traumatic Brain Injury, *Frontiers in Neurology* (2019). [DOI: 10.3389/fneur.2019.00825](#)

Provided by Kessler Foundation

Citation: Researchers explore neuromarkers for poor social outcomes after traumatic brain injury (2019, November 7) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-11-explore-neuromarkers-poor-social-outcomes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--