

Study raises new warnings about frail surgery patients

November 13 2019



Dr. Daniel Hall (left), seen here in a VA Pittsburgh Healthcare System operating suite with OR technician Mark Yost, is leading efforts to spread preoperative frailty screening in VA. Credit: Bill George

A Department of Veterans Affairs study has shown that frail surgery

patients may be at higher risk than previously thought. Mortality rates were high for frail patients even after surgeries normally considered low-risk. This led the researchers to conclude that "there are no 'low-risk' procedures among frail patients."

The findings appear in the Nov. 13, 2019, issue of *JAMA Surgery*.

Dr. Myrick C. Shinall, Jr. of the Vanderbilt University Medical Center, first author on the paper, explains that the findings change the view of which surgeries are considered dangerous. "We certainly found it surprising that surgeries usually considered low-stress and low-risk had such high risk of mortality for frail patients," he said in a phone interview. "We think other surgeons will also be surprised by the results."

Frailty refers to overall physical weakness. It is common in older adults. Frail patients have lowered resilience, energy, and ability to cope with stress. They show physical signs such as weakness and slowed activity.

As Shinall explains, the root cause of [frailty](#) is not completely known. Age is certainly a part of it, but not all people of a certain age become frail. Other co-occurring health problems, such as cardiovascular disease, play a part in patients becoming frail. Physical fitness also seems to protect against frailty to some extent.

While frailty is a well-known surgery concern, most research on frail patients has focused on high-risk surgeries. A surgery is usually considered high-risk when the post-surgery mortality rate is above 1%.

To test frailty's impact on surgeries generally considered less risky, the researchers looked at data on more than 400,000 veterans who had non-cardiac surgeries within the VA health care system. Out of this group, 8.5% were considered frail and 2.1% were very frail, based on a

measure called the Risk Analysis Index (RAI).

Patients were grouped based on what type of surgery they had. Low-risk surgeries included hernia surgery, appendectomy, and cyst removal. Surgeries such as amputation; arterial plaque removal; and knee, shoulder, or hip replacement were considered moderate-risk. High-risk procedures were mostly open surgeries on the aorta, lungs, liver or pancreas.

Mortality rates for frail and very frail patients were "alarmingly high" across all levels of surgery, compared to stronger patients. Thirty days after a low-risk surgery, 1.6% of frail patients and 10.3% of very frail patients had died. Non-frail patients had a 0.22% mortality rate after low-risk surgery.

For moderate-risk surgeries, 5.1% of frail patients and 18.7% of very frail patients died within 30 days. Mortality continued to rise for frail patients as time went on. The highest mortality rate recorded in the study was for very frail patients 180 days after moderate-risk surgery. About 43% of those patients had died.

Frailty rating remained a significant predictor of [mortality](#) regardless of the number of complications a patient had.

The results show that frailty adds significant risk to surgery, say the researchers, even when the surgery is considered minor.

Surgeons often do not consider whether patients can endure the stress of surgery for low-risk procedures, according to the researchers. Based on the results, the researchers recommend that patients be assessed using the RAI before any type of surgical procedure. If frailty is shown to be a risk, they say, surgeons could focus on pre-operative interventions to address it.

Assessing frailty could also help doctors and patients decide whether surgery is the best treatment option. As the researchers write in their report, "Even minor operations are high-risk for [frail patients](#); surgeons and referring physicians should consider if the potential benefits of [surgery](#) warrant the increased risk."

Dr. Daniel Hall, the corresponding author on the paper, is leading efforts to spread preoperative frailty screening in VA. The practice has been implemented in several VA medical centers already. Hall, a researcher at the VA Pittsburgh Healthcare System and the University of Pittsburgh, recently took part in a "Shark Tank" event held by VA's Diffusion of Excellence program. The VHA Shark Tank Competition offers VA employees the opportunities to share practices that address clinical and operational priorities to improve the veteran care experience. Through the competition, Hall received bids to implement the frailty screening at several more VA sites, earning Hall's practice the honor of "Gold Status."

Provided by Veterans Affairs Research Communications

Citation: Study raises new warnings about frail surgery patients (2019, November 13) retrieved 20 April 2024 from <https://medicalxpress.com/news/2019-11-frail-surgery-patients.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--