

Hospitalizations among dialysis patients are higher in areas with more black residents

November 10 2019

Patients with kidney failure who receive hemodialysis at US clinics located in residential areas with a high percentage of Black residents are more likely to be hospitalized than those who receive hemodialysis in communities with a lower percentage of Black residents. The findings come from a study that will be presented at ASN Kidney Week 2019 November 5-November 10 at the Walter E. Washington Convention Center in Washington, DC.

The analysis, conducted by Ladan Golestaneh MD, MS (Albert Einstein College of Medicine/Montefiore Medical Center) and her colleagues, included 4,567 patients on hemodialysis from 154 facilities in 127 zip codes in the United States. Patients receiving dialysis in communities with a high percentage of Black residents had higher hospitalization rates (32% higher for communities with 1.8% Blacks), despite having equivalent dialysis care. The association remained after adjusting for individual race, clinical comorbidities, community level poverty, and dialysis quality and adherence.

"Higher risk for hospitalization in communities with a high percentage of Blacks is likely not a result of differences in quality of dialysis care, adherence practices, or clinical factors, but rather a result of as yet unidentified community level determinants of health," said Dr. Golestaneh. "Healthcare disparities continue to exist even after measures taken by Centers for Medicare and Medicaid services to bridge the quality gap, through measures such as the dialysis Quality Incentive Program, to promote high-quality services in outpatient [dialysis](#) facilities

in [communities](#) across the U.S."

More information: Study: "Urban Segregation and Hospitalization Outcomes in Patients on Hemodialysis"

Provided by American Society of Nephrology

Citation: Hospitalizations among dialysis patients are higher in areas with more black residents (2019, November 10) retrieved 13 March 2024 from

<https://medicalxpress.com/news/2019-11-hospitalizations-dialysis-patients-higher-areas.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|