

# Kombucha, kimchi and yogurt: How fermented foods could be harmful to your health

November 13 2019, by Manal Mohammed

---



Credit: AI-generated image ([disclaimer](#))

Fermented foods have become very popular, thanks to claims about their nutritional properties and [reported health benefits](#), such as improving digestion, boosting immunity and even helping people lose weight. Some of the most popular fermented foods include kefir, kombucha,

sauerkraut, tempeh, natto, miso, kimchi and sourdough bread.

But though these fermented foods might offer us many health perks, most people aren't aware that they might not work for everyone. For some people, fermented foods might cause serious health issues.

Fermented foods are [loaded with microorganisms](#), such as live bacteria and yeast (known as probiotics). However, not all microorganisms are bad. Many, like probiotics, are harmless and are [even beneficial to us](#).

During the process of fermentation, probiotics convert carbohydrates (starch and sugar) into alcohol and/or acids. These act as a [natural preservative](#) and give fermented foods their distinctive zest and flavour. Many factors affect fermentation, including the type of [probiotic](#), the primary metabolites these microbes produce (such as lactic acid, or certain amino acids), and the [food](#) undergoing fermentation. For example, [probiotic yogurt](#) is produced by fermenting milk, most commonly with [lactic acid bacteria](#) which produce lactic acid.

Fermented foods contain high amounts of probiotics, which are generally considered safe for the majority of people. In fact, [they've been shown to have](#) anti-oxidant, anti-microbial, anti-fungal, anti-inflammatory, anti-diabetic and anti-atherosclerotic activity. However, some people might experience [severe side effects](#) after consuming fermented foods.

## 1. Bloating

The most common reaction to fermented foods is a [temporary increase in gas and bloating](#). This is the result of excess gas being produced after probiotics [kill harmful gut bacteria and fungi](#). Probiotics secrete antimicrobial peptides that kill harmful pathogenic organisms like Salmonella and E. Coli.

A recent study showed this [antimicrobial effect of probiotic Lactobacilli strains](#) found in commercial yogurt. Although bloating after eating probiotics seems to be a good sign that the harmful bacteria are being removed from the gut, some people might experience severe bloating, which can be very painful.

Drinking too much kombucha can also lead to excess sugar and calorie intake, which may also lead to [bloating and gas](#).

## 2. Headaches and migraines

Fermented foods rich in probiotics—including yogurt, sauerkraut and kimchi – [naturally contain biogenic amines](#) produced [during fermentation]. Amines are created by certain bacteria in order to [break down the amino acids](#) in fermented foods. The most common ones found in probiotic-rich foods include [histamine and tyramine](#).

Some people are sensitive to histamine and other amines, and [may experience headaches](#) after eating fermented foods. Because amines stimulate the central nervous system, they can increase or decrease blood flow, which can trigger headaches and migraines. One study found that low-histamine diets [reduced headaches](#) in 75% of participants. Taking a probiotic supplement might be therefore preferred.

## 3. Histamine intolerance

Histamine is [plentiful in fermented foods](#). For most, our body's specific enzymes will naturally digest them. However, some people don't produce enough of these enzymes. This means histamine won't be digested and will instead be absorbed into the bloodstream.

This can cause a range of [histamine intolerance symptoms](#). The most

common are itching, headaches or migraines, runny nose (rhinitis), eye redness, fatigue, hives and digestive symptoms include diarrhoea, nausea and vomiting.

However, histamine intolerance can also cause more severe symptoms, including asthma, low blood pressure, irregular heart rate, circulatory collapse, sudden psychological changes (such as anxiety, aggressiveness, dizziness and lack of concentration) and sleep disorders.

## **4. Food-borne illness**

While most fermented foods are safe, it's still possible for them to get contaminated with bacteria that can cause illness. In 2012, there was an outbreak of 89 cases of Salmonella in the US because of [unpasteurised tempeh](#).

Two large outbreaks of Escherichia coli, were reported in South Korean schools in 2013 and 2014. These were associated with consumption of [contaminated fermented vegetable kimchi](#).

In most cases, probiotics found in fermented [milk products](#) such as cheese, yogurt and buttermilk can effectively prevent the growth of certain bacteria, such as [Staphylococcus aureus](#) and Staphylococcal enterotoxins which can cause food poisoning. But in some cases probiotics fail and bacteria can actually secrete toxins, so the product may be hazardous.

## **5. Infection from probiotics**

Probiotics are generally safe for the vast majority of people. However, in rare cases, they can cause infection—especially in people who [have a compromised immune system](#).

A London study reported the first case of a 65-year-old diabetic patient whose liver abscess had been [caused by probiotic consumption](#).

Susceptible patients, such as those with compromised immunity, should be advised against excessive consumption of probiotics.

Treatment with probiotics [can cause serious infections](#) such as pneumonia in vulnerable people, as well as systemic infections including [sepsis](#) and [endocarditis](#).

## 6. Antibiotic resistance

Probiotic bacteria can carry genes that confer resistance to antibiotics. These antibiotic resistance genes may pass to other bacteria found in the food chain and gastrointestinal tract via [horizontal gene transfer](#). The most common [antibiotic resistance](#) genes carried by fermented foods are against [erythromycin and tetracycline](#), which are used to treat respiratory infections and some sexually transmitted diseases.

Researchers found resistant probiotic strains in commercially available dietary supplements, which could mean resistance to several common types of antibiotics used to treat [serious bacterial infections](#).

Research has also found six probiotic *Bacillus* strains found in food products (including kimchi, yogurt and olives) are also [resistant to several antibiotics](#).

And, a recent Malaysian study showed probiotic [Lactobacilli bacteria](#) in kefir carry resistance to numerous antibiotics, including ampicillin, penicillin and tetracycline. These are used to treat serious human diseases including bladder infections, pneumonia, gonorrhea, and meningitis.

Another study also showed lactic acid [bacteria](#) found in Turkish dairy

products were [resistant mainly to vancomycin antibiotic](#), which is the drug of choice for treatment of [MRSA infection](#).

While there are a wide variety of [health benefits](#) that can happen from consuming fermented foods, these may not work for everyone. While most people will be fine eating fermented foods, for some they could potentially cause serious health problems.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Kombucha, kimchi and yogurt: How fermented foods could be harmful to your health (2019, November 13) retrieved 4 May 2024 from <https://medicalxpress.com/news/2019-11-kombucha-kimchi-yogurt-fermented-foods.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--