

Mapping the pathway to gut health in HIV and SIV infections

November 19 2019



Satya Dandekar and Katti Crakes from UC Davis Health. Credit: UC Davis Health

A UC Davis study found that the damaged gut lining (known as leaky gut) in monkeys infected with chronic simian immunodeficiency virus



(SIV), an HIV-like virus, was rapidly repaired within five hours of receiving *Lactobacillus plantarum* bacteria.

The study, published today in the *PNAS*, linked chronically inflamed <u>leaky gut</u> to the loss of PPAR α signaling (a nuclear receptor protein responsible for regulating <u>cell metabolism</u>) and subsequent damage to mitochondria—the cell's power house.

The researchers found that *L. plantarum* activated PPAR α signaling and revived mitochondrial flow, repairing the gut barrier in only five hours of exposure.

The outcome lends hope that leaky gut, a common condition among HIV patients, could be effectively treated in the future.

HIV and the damage to the gut lining

The gut, home to majority of the lymphoid tissue in the body, is an early target of HIV. The virus severely damages the immune and epithelial cells in the gut's lining. This damage leads to an inflamed and leaky gut with weakened defense system and decreased nutrient absorption.

Anti-retroviral therapy (ART) for HIV has been successful in limiting the damage to the body's <u>immune system</u>. Yet, it has failed to consistently or completely repair the damage to the gut and its lining.

"We wanted to map the pathways that lead to sustained damage in the gut and to identify ways to intervene and support its repair," said Katti Crakes, doctoral student in the schools of Medicine and Veterinary Medicine at UC Davis and first author on the study.

The researchers found that HIV attacks the gut's epithelial cells by targeting and draining the mitochondria.



"To reverse the damage caused by HIV and to increase the efficacy of ART, it is important to restore mitochondrial function and to rapidly repair the gut epithelium and immune defense," said Satya Dandekar, professor of microbiology and immunology at UC Davis School of Medicine and senior author.

Identifying cell signaling regulators for restoring the gut barrier

The bacteria present in the gut are known to play an important role in supporting and repairing the gut functioning. The study specifically tested the impact of *L. plantarum* bacteria on gut epithelial barrier of SIV infected rhesus macaques.

"We challenged the capacity of *L. plantarum* bacteria and their metabolites to restore the gut functions in an extremely inflamed visibly disrupted gut environment," Dandekar said.

The researchers found that *L. plantarum* were able to survive and remain metabolically active in inflamed gut. The bacteria repaired the gut barrier by targeting and restoring the mitochondria in the intestinal <u>epithelial cells</u> damaged by SIV as well as HIV. These findings provide translational insights into restoring gut immunity and function, both of which are essential for successful HIV cure efforts.

More information: Katti R. Crakes et al, PPARα-targeted mitochondrial bioenergetics mediate repair of intestinal barriers at the host–microbe intersection during SIV infection, *Proceedings of the National Academy of Sciences* (2019). DOI: 10.1073/pnas.1908977116



Provided by UC Davis

Citation: Mapping the pathway to gut health in HIV and SIV infections (2019, November 19) retrieved 5 May 2024 from https://medicalxpress.com/news/2019-11-pathway-gut-health-hiv-siv.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.