

# Among people with bipolar disorder, inflammation predicts cognitive deficits

November 19 2019

---



Bipolar disorder is characterized by transitions between depression and mania.  
Credit: Wikipedia

Bipolar disorder is a heterogeneous disease—two people who share the

same diagnosis can have very different outcomes and quality of life. Treatment of bipolar disorder largely focuses on controlling episodes of depression or mania, but even when these symptoms are well managed, some patients will suffer from cognitive deficits while others will show cognitive resilience.

Understanding why people with the same clinical diagnosis have such different symptoms and outcomes could lead the way to better treatments. A team of investigators from Brigham and Women's Hospital has conducted the largest study to date of people with [bipolar disorder](#) that examined whether [inflammation](#) may play an important role in patient outcomes. The team reports that levels of a biomarker of inflammation were significantly higher among patients with poor cognitive performance, pointing to a potentially modifiable risk factor for cognitive deficits. Their findings are published in *Molecular Psychiatry*.

"We know that one person with bipolar disorder may be very different from another, and these findings support this. Levels of inflammation varied in patients, but among those with high levels of inflammation, we saw the strongest evidence of cognitive impairment," said corresponding author Katherine Burdick, Ph.D., the associate vice chair for Research in the Department of Psychiatry and director of the Mood and Psychosis Research Program at the Brigham. "Hundreds of things can contribute to cognitive decline, from poor sleep to lack of exercise to poor diet. There's mounting evidence that inflammation may be an important driver, too."

To conduct their study, Burdick, lead author Caitlin Millett, Ph.D., a postdoctoral research fellow in the Mood and Psychosis Research Program, and their colleagues recruited patients with bipolar disorder whose mood was clinically stable at the time of assessment. The study included 222 participants with bipolar disorder and compared their

results to 52 healthy controls. Participants underwent a battery of cognitive tests, and the team measured levels of C-reactive protein (CRP), a marker of systemic inflammation and risk of inflammatory disease.

The team found that levels of the biomarker for inflammation were tied to cognitive function. High levels of CRP significantly predicted cognitive decline among participants with bipolar disorder, even after controlling for factors such as age, education and more.

Previous studies by other Brigham investigators built the case that high levels of CRP are both a marker of inflammation and a predictor of heart attack risk. Inflammation and CRP are now tied to risk of many diseases—some of them unexpected—including colon cancer, obstructive sleep apnea, rheumatoid arthritis and more. Burdick, Millett and colleagues note that because CRP is such a global marker of inflammation, further study of patients with bipolar disorder is needed. The team already has plans in place to conduct a longitudinal study of the participants, repeatedly measuring CRP levels and other more specific immune markers to understand how they change over time. This will provide a clearer picture of whether inflammation causes [cognitive decline](#).

"Cognitive impairment is a core component of bipolar disorder. Because of its significant role in predicting a person's level of functioning in their everyday lives, this needs to be considered a treatment target," said Burdick. "The message here is an optimistic one: This is a marker of something we can actively change, which opens the door for hope."

Provided by Brigham and Women's Hospital

Citation: Among people with bipolar disorder, inflammation predicts cognitive deficits (2019,

November 19) retrieved 25 April 2024 from <https://medicalxpress.com/news/2019-11-people-bipolar-disorder-inflammation-cognitive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.