


Researchers find potential new target for treatment of inflammatory disease

November 7 2019



Credit: CC0 Public Domain

Researchers led by the University of Birmingham have found a potential new target to treat inflammatory disease.

The research, led by scientists at the University of Birmingham's Institute of Inflammation and Ageing, Institute of Cardiovascular Sciences, and Institute of Metabolism and Systems Research, publishes



today in Cell Metabolism.

The team of researchers are experts in cellular metabolism—the set of chemical reactions or 'metabolic pathways' that occur in living organisms in order to maintain life.

This study focussed on the role of <u>lactate</u> in how the body's <u>immune</u> <u>system</u> responds to inflammation. Lactate is a molecule produced by most tissues in the human body, with the highest production found in muscle, and levels of lactate in the blood are regulated by the kidney.

A high lactate level in the blood means that a disease or condition is causing lactate to accumulate. In general, a greater increase in lactate means a greater severity of the condition, and lactate levels are used by hospital medics as a critical way to monitor the health and likely recovery of patients in <u>intensive care</u>.

Dr. Valentina Pucino, of the University of Birmingham, said: "The recent discovery of the fundamental role of metabolism in immune cell biology is contributing immensely to our understanding of immune cell regulation."

Dr. Michelangelo Certo, of the University of Birmingham, added: "So far, most studies have focused on the role of metabolic pathways in the establishment of the immune response.

"Lactate has mainly been seen as a by-product of metabolism or as a biomarker in critical care at best, rather than a bioactive molecule, and its functional effects have thus been neglected for a long time.

"However, far from being inert, the accumulation of lactate in the cells, molecules and structures of tissue is found in both inflammatory disease and cancer."



In this study, the researchers analysed the response of immune cells to lactate in <u>chronic inflammation</u> caused by <u>rheumatoid arthritis</u>.

In their research, which involved the use of mice, they also used blood from healthy patients and patients with arthritis, as well as joint biopsies, and tonsils removed following tonsillectomies due to the fact that inflamed tonsils share similarities with an inflamed joint.

Dr. Claudio Mauro, of the University of Birmingham's Institute of Inflammation and Ageing, said: "We identified the pathway initiated by lactate build-up in inflamed tissue that exacerbates the inflammatory response.

"We also now can provide evidence that molecules made by immune <u>cells</u> to transport sodium lactate—the sodium salt of lactic acid—could be a target for treatment to stop this lactate build-up in chronic inflammatory disorders."

More information: Pucino et al (2019). 'Lactate build-up at the site of chronic inflammation promotes disease by inducing CD4+ 1 T cell metabolic rewiring.' *Cell Metabolism*.

Provided by University of Birmingham

Citation: Researchers find potential new target for treatment of inflammatory disease (2019, November 7) retrieved 1 May 2024 from https://medicalxpress.com/news/2019-11-potential-treatment-inflammatory-disease.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.