


Reversed halo signs manifest in septic pulmonary embolism due to IV drug use

November 1 2019



Axial lung window chest CT image shows two reversed halo signs in anterior segment (arrows) of right upper lobe with central reticulation. Credit: American Journal of Roentgenology (AJR)

According to an article published ahead-of-print in the January 2020

issue of the *American Journal of Roentgenology (AJR)*, the reversed halo sign was frequently observed on the chest CT scans of patients with IV substance use disorder-related septic pulmonary embolism (PE).

Of the 62 patients (54.8% women; 32.8 ± 8.3 [SD] years) who met Harvard Medical School radiologist Renata R. Almeida and colleagues' inclusion criteria—IV substance use disorder, findings of septic PE on chest CT scans, and confirmation of infection—59.7% (37/62) had reversed [halo](#) signs ($\kappa = 0.837-0.958$, p

Moreover, the mean number of unique reversed halo signs per patient was 2.1 ± 1.7 , with 46.7% of patients having more than one reversed halo sign.

Noting that the reversed halo sign was an early and reliable imaging finding observed in most cases of CT-based diagnosis of septic PE secondary to IV substance use disorder, as Almeida et al. concluded, "septic PE should be included in the differential diagnosis of patients presenting with the reversed halo sign and history of IV substance use disorder."

The authors added: "Apparent differences of frequency, shape, and distribution in comparison with case series of pulmonary thromboembolism and invasive fungal infections could be investigated by future studies as possible biomarkers to assist in discrimination between septic PE and other causes of pulmonary infarct manifesting with the reversed halo sign."

More information: Renata R. Almeida et al, Frequency and Reliability of the Reversed Halo Sign in Patients With Septic Pulmonary Embolism Due to IV Substance Use Disorder, *American Journal of Roentgenology* (2019). [DOI: 10.2214/AJR.19.21659](https://doi.org/10.2214/AJR.19.21659)

Provided by American Roentgen Ray Society

Citation: Reversed halo signs manifest in septic pulmonary embolism due to IV drug use (2019, November 1) retrieved 11 May 2024 from <https://medicalxpress.com/news/2019-11-reversed-halo-manifest-septic-pulmonary.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.