


# Sex workers' preferences for HIV prevention center on convenience

November 13 2019, by Misti Crane

---



Microscopic image of an HIV-infected T cell. Credit: NIAID

Preventing HIV in sex workers is a powerful tool in lowering the worldwide burden of the disease, and a new study could help ensure that high-risk women take advantage of medical safeguards.

Researchers studying HIV in Malawi knew that the drug PrEP (pre-exposure prophylaxis) could prevent infections among [female sex workers](#) and their [sex partners](#), but they wanted to know how best to get them the medication and encourage them to use it. So they asked the [women](#) themselves.

The women in the study said that the most important considerations for them were the location where they'd receive the medication and the additional services they could receive during those appointments. In particular, they were interested in cervical cancer screenings.

In terms of location, the majority of sex workers preferred to receive the HIV-prevention medication at family planning clinics or non-governmental drop-in centers—not at mobile health clinics near where they live or work.

"We really need to think about integrating care for women, providing them these services at locations where they already feel comfortable and can receive other essential health care," said lead researcher Kathryn Lancaster, an assistant professor in Ohio State's College of Public Health. The study appears today (Nov. 13, 2019) in the journal *AIDS and Behavior*.

PrEP, a daily oral regimen of medication, reduces the risk of getting HIV from sex by about 99 percent when taken as recommended, according to the Centers for Disease Control and Prevention.

"In Malawi, HIV infections have been on the decline in recent years with better education, testing and access to prevention. Nationally, about 10 percent of the population is infected, but about 70 percent of women who are sex workers are currently infected," Lancaster said.

Barriers to treatment for active infections and to prevention for those

who aren't already HIV-positive are significant for those who are poor and working in a stigmatized, yet legal, field in Malawi, she said.

"There's an inherent imbalance in power in these transactions. And the women often find themselves forced into sex or may be paid more for having unprotected sex," Lancaster said.

That makes PrEP an especially important tool because it is something women can use to protect themselves without their sexual partners' cooperation. And more PrEP use among the workers would also mean fewer infected male customers.

In March of this year, the Malawi government OK'd use of PrEP specifically for sex workers, but there are no guidelines on how to make that happen, Lancaster said.

She's hopeful that the information collected in this new study, which included 44 sex workers in eight focus groups, will guide programs aimed at preventing HIV infection among these women.

"Convenience was the main underlying factor for the sex workers' preferences and it is important for the government and others interested in prevention to understand how to best serve the needs of the women they hope to protect," Lancaster said.

"Currently, the best data we have shows that just a few hundred Malawi [sex workers](#) at best are on PrEP when there are thousands who could benefit. There's so much good that could come of providing PrEP to these women, but if we don't make it convenient for them, they're not going to use it," Lancaster said.

For instance, providing prevention care at a site that is specific to treatment for sexually transmitted infections could be a deterrent, she

said.

"They preferred family planning clinics and I think that speaks to feeling more comfortable with receiving this care alongside a service they're already using," she said.

Along with convenient locations and extra health services, the women also said they preferred male health care providers and a setup where they would receive two months' worth of PrEP at a time.

**More information:** Kathryn E. Lancaster et al. Preferences for Pre-exposure Prophylaxis Service Delivery Among Female Sex Workers in Malawi: A Discrete Choice Experiment, *AIDS and Behavior* (2019).

[DOI: 10.1007/s10461-019-02705-3](https://doi.org/10.1007/s10461-019-02705-3)

Provided by The Ohio State University

Citation: Sex workers' preferences for HIV prevention center on convenience (2019, November 13) retrieved 20 April 2024 from <https://medicalxpress.com/news/2019-11-sex-workers-hiv-center-convenience.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.