

Stress, plastic additives in late pregnancy raise risk of premature birth

November 14 2019, by Patti Verbanas



Mothers in the study who had higher levels of phthalates in their urine and reported stress during their pregnancy as well were more likely to give birth preterm. Credit: Rutgers University

Women exposed simultaneously to stress and plastic additives late in pregnancy are at increased risk for premature birth, according to a study by Rutgers and other institutions.



The study, published in the journal *Environment International*, is the first to analyze a link between stress and phthalates—a group of chemicals in plastics, <u>personal care products</u> and electronics—and premature births. The findings are the latest in the Infant Development and the Environment Study, which tracked 783 women throughout their pregnancies between 2010 and 2012.

"Both exposure to phthalates and high levels of stress have been individually linked to births before 37 weeks gestation, but how these two <u>risk factors</u> may influence each other had not been previously explored," said co-author Emily Barrett, an associate professor at Rutgers School of Public Health and Environmental and Occupational Health Sciences Institute. "Our research suggests that the third trimester is the critical window for these risks."

People are exposed to phthalates by eating and drinking foods from containers and products containing the chemicals, through packaging and food handling equipment, through their skin in personal care products and in the air they breathe.

While stress itself is not related to the exposure of these <u>plastic additives</u>, stress can affect the <u>immune system</u>, lead to inflammation and change hormone levels, potentially making these women more vulnerable to the adverse effects of phthalates.

Barrett said other factors—how a woman perceives stress, how she uses social supports to lessen stress and the total number of stress events in her life—can affect how stress during pregnancy affects her child's health. The researcher recommends that pregnant women limit their exposure to phthalates by eating less processed foods and reducing use of plastics and personal care products.

The researchers analyzed urine samples throughout the women's



pregnancies for traces of phthalates and reviewed questionnaires they completed in their third trimester about stressful events—such as job loss, serious illness, family death, relationship difficulties and legal or financial problems—during their pregnancies. About nine percent of the women delivered prematurely, 70 percent of which were spontaneous.

"Mothers who had higher levels of phthalates in their urine and reported stress during their pregnancy as well were more likely to give birth preterm," Barrett said. "We think that <u>stress</u> may make the body more vulnerable to the impacts of chemicals in our environment, like phthalates."

More information: Kelly K. Ferguson et al, Joint impact of phthalate exposure and stressful life events in pregnancy on preterm birth, *Environment International* (2019). DOI: 10.1016/j.envint.2019.105254

Provided by Rutgers University

Citation: Stress, plastic additives in late pregnancy raise risk of premature birth (2019, November 14) retrieved 30 April 2024 from <u>https://medicalxpress.com/news/2019-11-stress-plastic-additives-late-pregnancy.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.