


Many teens have low iron, B12 levels years after bariatric surgery

November 8 2019



Prevalence of 2 or more nutritional deficiencies increased significantly over 5 years in adolescents after gastric bypass, but not after sleeve gastrectomy. Image adapted from Supplemental Figure 1 from Xanthakos et al. *Clinical Gastroenterology and Hepatology* 2019 (in press)

This chart shows that vertical sleeve procedures pose fewer long-term nutritional disruptions for teens who need treatment for severe obesity. More details from this study led by Cincinnati Children's appear in *Clinical Gastroenterology and Hepatology*. Credit: Cincinnati Children's

The good news: bariatric surgery can be a life-changer for severely obese teens. Not only do young people experience dramatic weight loss, they also benefit from near-immediate improvement of diabetes plus longer-term reductions in heart health risks.

However, these benefits come with a caution: recipients may develop [nutritional deficiencies](#) years after treatment that can carry their own health risks if not properly managed. Also, one of the two primary forms of [bariatric surgery](#), [gastric bypass](#), appears to cause more nutritional disruption than the other, vertical sleeve gastrectomy.

Findings from this study, led by experts at Cincinnati Children's, were published online Nov. 8, 2019 in the journal *Clinical Gastroenterology and Hepatology*.

"This study highlights the importance of annual nutritional screening", says Stavra Xanthakos, MD, Medical Director, Surgical Weight Loss Program for Teens at Cincinnati Children's and first author of the study. "We also found that the risk of nutritional deficiency is lowest in patients with the highest intake of recommended vitamin and mineral supplements, so these risks can be minimized with appropriate supplementation and good adherence."

In this study, Xanthakos, senior author Heidi Kalkwarf, Ph.D., RD and their surgical colleagues, including Michael Helmuth, MD at Cincinnati Children's and Thomas Inge, MD, Ph.D. at Children's Hospital Colorado, focused on the nutritional impacts of the two most common forms of the [surgery](#): Roux-en-Y gastric bypass (RYGB) and vertical sleeve gastrectomy (VSG). Their findings, five years after treatment:

- In both procedures, patients benefitted from weight loss of approximately 23%
- After RYGB, levels of vitamin B12 declined significantly, but

B12 levels did not drop after VSG.

- After RYGB, 71% of patients had low iron stores compared to 2.5% of patients prior to surgery.
- For those receiving VSG, 45% had low iron stores at five years, compared to 11% prior to surgery.
- Over twice as many RYGB participants than VSG recipients had multiple nutritional deficiencies at 5 years (59% vs. 27%).
- No [significant changes](#) were found after either procedure for folate or vitamins A, B1, or D.

What are the risks from low iron and low B12?

"Both B12 and [iron deficiency](#) can cause anemia, and B12 deficiency can also cause significant neurological dysfunction," says senior author Kalkwarf. "If caught early, [dietary supplements](#) can minimize these risks, which is why annual nutrition testing is so important. It's also important to eat a well-balanced diet in addition to taking supplements and involve a registered dietitian in the pre- and post-operative care of these patients"

Vertical Sleeve Gastrectomy



Roux-en-Y Gastric Bypass



Data from the national Teen-LABS study suggests that vertical sleeve procedures should be the first-line option when teens with severe obesity need surgical intervention. Credit: Cincinnati Children's

Untreated, the most common symptoms of iron deficiency anemia in adults include fatigue, weakness, and exercise intolerance. In young children, iron deficiency also can lead to cognitive impairment, but it is not known if developing iron deficiency in adolescence would have the same effect.

People with B12 deficiency also can suffer from anemia, but also can

experience symptoms of depression, irritability, cognitive slowing, and forgetfulness. With severe deficiency, some can develop numbness or tingling in extremities, gait problems, and partial paralysis.

Benefits of surgery outweigh these risks

"In our bariatric surgery program at Cincinnati Children's, all patients get annual micronutrient and nutritional assessments as recommended. It's also important for them to continue annual nutritional screening after they transition to adult care later in life," Xanthakos says.

"The positive message is that the risk of nutritional deficiency can be minimized with appropriate dietary supplementation, so these procedures should not be avoided solely due to nutritional concerns," says Inge, Teen-LABS Principal Investigator. "These procedures are very helpful for severely obese adolescents, especially those struggling with diabetes, severe fatty liver disease or sleep apnea."

Longer term follow-up is also important to detect any other consequences from surgery, such as changes in bone health. "For pediatric surgeons, these findings suggest favoring the VSG procedure as the first line approach in teens," says Inge.

Meanwhile, newer endoscopic bariatric procedures, approved for adults may eventually become available at younger ages. "We will need to study nutritional risks after these endoscopic procedures as well," Xanthakos says.

More information: Stavra A. Xanthakos et al, Nutritional Risks in Adolescents After Bariatric Surgery, *Clinical Gastroenterology and Hepatology* (2019). [DOI: 10.1016/j.cgh.2019.10.048](https://doi.org/10.1016/j.cgh.2019.10.048)

Provided by Cincinnati Children's Hospital Medical Center

Citation: Many teens have low iron, B12 levels years after bariatric surgery (2019, November 8)
retrieved 2 May 2024 from

<https://medicalxpress.com/news/2019-11-teens-iron-b12-years-bariatric.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.