

15-year study finds treatment gaps exist for patients with peripheral arterial disease

November 18 2019



New 15-year study found that patients with peripheral arterial disease may not be prescribed these life-saving medications at the same rate as for other atherosclerotic conditions, such as heart attack or stroke. Credit: Intermountain Healthcare

Peripheral arterial disease (PAD) is a progressive and life-threatening

form of atherosclerosis that narrows the arteries serving the legs, arms, chest, and abdominal organs. It affects more than 8.5 million people in the United States, including 10 to 20 percent of individuals over 60 years old.

Outcomes of PAD patients are worse than those with [coronary disease](#) or [cerebrovascular disease](#), emphasizing the importance of attention to their treatment. Like other kinds of atherosclerotic cardiovascular diseases, it should be treated with high-intensity doses of statins.

However, a new 15-year study by researchers at the Intermountain Healthcare Heart Institute in Salt Lake City found that PAD patients may not be prescribed these life-saving medications at the same rate as for other atherosclerotic conditions, such as [heart attack](#) or stroke.

"As a profession we focus so much on coronary disease that we may be not be optimally treating patients with PAD, especially when it comes to statin prescription," said Jeffrey L. Anderson, principal investigator of the study and distinguished clinical and research physician at the Intermountain Healthcare Heart Institute. "However, we found that over time, statin prescription rates across our healthcare system have increased for PAD patients, and with it, PAD patient outcomes have improved. Despite this, there remains a significant treatment gap, so much more progress is needed."

In the study, researchers identified 4,333 patients who had a first encounter with PAD (disease in the aorta and lower extremities) between January 1, 1999 and December 31, 2013, and survived that event.

They then were followed for at least three years or until their death. Overall, statin prescription was associated with a reduced 3-year rate of subsequent [heart](#) attack, stroke, or death, from 23.5% to 18.9%, and it

was only 14.3% if the prescription was for a high-intensity statin. Unfortunately, only one-third of patients overall had been prescribed a statin at discharge after their PAD event, and only 11% were prescribed a high-intensity statin.

The Intermountain researchers then gave attention to the treatment trend over 15 years. From 1999 through 2013, a statin prescription increased in five-year increments from 17% to 37% to 48%, indicating steady progress toward the 100% ultimate treatment goal. However, the most recent value of 48% for PAD still lags significantly behind values of 74% for cerebrovascular disease (e.g., stroke) patients and 93% for coronary [disease](#) (e.g., heart attack) patients.

Findings from the study will be presented at the 2019 American Heart Association Scientific Sessions Philadelphia on Nov. 17, 2019.

Dr. Anderson noted the favorable progress in PAD treatment, but he highlighted the remaining large treatment gap when compared to coronary and stroke patients, showing that more work needs to be done in statin prescribing for patients with PAD.

"We're doing much better in terms of treating these patients with statins, but there's still a need for more attention to this group," he said. "More than 90 percent of our heart attack patients are prescribed statins. We need to keep focusing on PAD patients to get them to the right physicians who are fully aware of the need to treat them as high-risk patients and to get their statin percentages also up into the mid-90s. These patients represent the highest risk atherosclerosis group, but they also are the group that stands to benefit the most from [statin](#) therapy."

Provided by Intermountain Medical Center

Citation: 15-year study finds treatment gaps exist for patients with peripheral arterial disease (2019, November 18) retrieved 6 May 2024 from <https://medicalxpress.com/news/2019-11-year-treatment-gaps-patients-peripheral.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.