


## Study finds BPA levels in humans dramatically underestimated

December 5 2019



3D chemical structure of bisphenol A. Credit: Edgar181 via Wikimedia Commons

Researchers have developed a more accurate method of measuring bispehnol A (BPA) levels in humans and found that exposure to the endocrine-disrupting chemical is far higher than previously assumed.



The study, published in the journal the *Lancet Diabetes & Endocrinology* on Dec. 5, provides the first evidence that the measurements relied upon by <u>regulatory agencies</u>, including the U.S. Food and Drug Administration, are flawed, underestimating <u>exposure</u> levels by as much as 44 times.

"This study raises serious concerns about whether we've been careful enough about the safety of this chemical," said Patricia Hunt, Washington State University professor and corresponding author on the paper. "What it comes down to is that the conclusions <u>federal agencies</u> have come to about how to regulate BPA may have been based on inaccurate measurements."

BPA can be found in a wide range of plastics, including food and drink containers, and animal studies have shown that it can interfere with the body's hormones. In particular, fetal exposure to BPA has been linked to problems with growth, metabolism, behavior, fertility and even greater cancer risk.

Despite this experimental evidence, the FDA has evaluated data from studies measuring BPA in human urine and determined that human exposure to the chemical is at very low, and therefore, safe levels. This paper challenges that assumption and raises questions about other chemicals, including BPA replacements, that are also assessed using indirect methods.

Hunt's colleague, Roy Gerona, assistant professor at University of California, San Francisco, developed a direct way of measuring BPA that more accurately accounts for BPA metabolites, the compounds that are created as the chemical passes through the human body.

Previously, most studies had to rely on an indirect process to measure BPA metabolites, using an enzyme solution made from a snail to



transform the metabolites back into whole BPA, which could then be measured.

Gerona's new method is able to directly measure the BPA metabolites themselves without using the enzyme solution.

In this study, a research team comprised of Gerona, Hunt and Fredrick vom Saal of University of Missouri compared the two methods, first with synthetic urine spiked with BPA and then with 39 human samples. They found much higher levels of BPA using the direct method, as much as 44 times the mean reported by the National Health and Nutrition Examination Survey (NHANES). The disparity between the two methods increased with more BPA exposure: the greater the exposure the more the previous method missed.

Gerona, the first author on the paper, said more replication is needed.

"I hope this study will bring attention to the methodology used to measure BPA, and that other experts and labs will take a closer look at and assess independently what is happening," he said.

The research team is conducting further experiments into BPA measurement as well as other chemicals that may also have been measured in this manner, a category that includes environmental phenols such as parabens, benzophenone, triclosan found in some cosmetics and soaps, and phthalates found in many <u>consumer products</u> including toys, food packaging and <u>personal care products</u>.

"BPA is still being measured indirectly through NHANES, and it's not the only endocrine-disrupting <u>chemical</u> being measured this way," Gerona said. "Our hypothesis now is that if this is true for BPA, it could be true for all the other chemicals that are measured indirectly."



## **More information:** *Lancet Diabetes & Endocrinology*, <u>DOI:</u> <u>10.1016/S2213-8587(19)30381-X</u>

## Provided by Washington State University

Citation: Study finds BPA levels in humans dramatically underestimated (2019, December 5) retrieved 13 May 2024 from <a href="https://medicalxpress.com/news/2019-12-bpa-humans-underestimated.html">https://medicalxpress.com/news/2019-12-bpa-humans-underestimated.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.