

# Breast cancer patients with government insurance at higher risk of death

December 13 2019

---



Samilia Obeng-Gyasi, MD (The Ohio State University) is the lead author on an analysis of the insurance status of 9800 breast cancer patients and their clinical outcomes. The analysis was conducted through the ECOG-ACRIN Cancer Research Group. Credit: ECOG-ACRIN Cancer Research Group

A retrospective study of nearly 9800 women with breast cancer who participated in randomized clinical trials was presented today at the 2019 San Antonio Breast Cancer Symposium. The study found that women with government insurance (Medicaid or Medicare) were much less likely to participate in a clinical trial compared to their privately insured counterparts. The few women with government insurance who did participate in the trials were more likely to stop treatment early and had lower survival. The study was conducted by researchers in the ECOG-ACRIN Cancer Research Group (ECOG-ACRIN) with funding from the

National Cancer Institute, part of the National Institutes of Health.

"Having insurance in and of itself was not enough to ensure access to a clinical trial or a good clinical outcome," said lead researcher Samilia Obeng-Gyasi, MD, a breast surgical oncologist at The Ohio State University who studies social determinants of cancer care.

Dr. Obeng-Gyasi and colleagues looked at data from women who participated in two large ECOG-ACRIN trials testing chemotherapy treatments. Study E1199 enrolled participants from 1999 to 2002 and study E5103 enrolled participants between 2007 and 2011. In the current analysis, Dr. Obeng-Gyasi and colleagues evaluated each participant's [insurance coverage](#) and neighborhood [socioeconomic status](#) at the time they entered the trial, to determine if either status affected the women's clinical outcomes later on.

Insurance status was either private, government (Medicaid, Medicare), or self-pay. Only about 13% of participants in each of the two [trials](#) had [government insurance](#). The majority of patients had [private insurance](#) (85.6% in E1199 and 82% in E5103).

Patients with government insurance were less likely to complete treatment compared to their privately insured counterparts. One out of every four government insured patients did not complete treatment in E1199 versus one out of every seven privately insured women. In E5103, approximately one out of every two government insured patients did not complete therapy compared to one out of every three privately insured patients. Completion of the trial and survival outcomes were each controlled for disease severity, age, and other patient characteristics that could influence the participant's outcomes.

The government insured patients also had a higher risk of dying compared to their privately insured counterparts. In E1199, the risk of

death increased by one half. For those in E5103, the risk increased by one-third.

The researchers determined the neighborhood socioeconomic status of each participant by linking their zip code to county-level data on occupation, income, poverty, wealth, education, and crowding.

There was no association between neighborhood socioeconomic status and trial completion or survival.

"Insurance appears to be a powerful determinant of clinical trial outcomes," said Dr. Obeng-Gyasi. "With continued changes to insurance at the federal and state levels, physicians and policy-makers would benefit from having more data such as what we produced in our study. Right now, most [clinical trials](#) do not collect data on social determinants of health. If they did, the information could lead us to better understand the interaction of insurance status and many other factors on the clinical outcomes of clinical trial participants."

**More information:** [Abstract PD10-09](#): Impact of insurance and socioeconomic status on clinical outcomes in therapeutic clinical trials for breast cancer. Category: Disparities and Barriers to Care. First author: Samilia Obeng-Gyasi. PD10 Spotlight Session: Toxicity, Tolerability & Cost (Friday, December 13, 7:00 AM.)

Provided by ECOG-ACRIN Cancer Research Group

Citation: Breast cancer patients with government insurance at higher risk of death (2019, December 13) retrieved 18 April 2024 from <https://medicalxpress.com/news/2019-12-breast-cancer-patients-higher-death.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.