

# Bystanders can help more cardiac arrest victims survive

December 5 2019

---



Only 8% of Americans survive cardiac arrest outside a hospital, but that percentage could increase significantly if bystanders recognize cardiac arrest and perform simple lifesaving tasks, says UVA Health physician William J. Brady, MD. Credit: UVA Health

Only 8% of Americans survive cardiac arrest outside a hospital, but that percentage could increase significantly if bystanders recognize cardiac

arrest and perform simple lifesaving tasks, a UVA Health physician says.

UVA emergency medicine physician William J. Brady, MD, co-authored an article in the *New England Journal of Medicine* that highlights a few steps anyone without extensive medical training can do to help:

- Recognize quickly the signs of cardiac arrest and call 911
- Begin cardiopulmonary resuscitation (CPR) using chest compressions
- Use an automated external defibrillator (AED), when appropriate, to deliver an electric current to restart a victim's heart

"Early emergency care by the [bystander](#), long before emergency medical providers arrive, can save a life. This care is referred to as 'pre-arrival care,'" said Brady. "This pre-arrival care is as important as more advanced care by EMTs, nurses, and doctors once trained providers are available."

## Calling 911, Beginning CPR

The first step is simply noting that someone may be in cardiac arrest.

The article notes that the American Heart Association and other medical groups "recommend that bystanders should suspect cardiac arrest and begin CPR whenever a person is unresponsive and not breathing normally." Faster recognition of cardiac arrest will also lead to a quicker 911 call, speeding the arrival of trained emergency medical providers.

But research has shown that bystanders can help cardiac arrest victims by beginning CPR themselves (major groups like the American Heart Association now endorse compression-only CPR, which is performed using chest compressions only and not breathing into the victim's

mouth). A 2010 review of more than 10,000 out-of-hospital cardiac arrest cases found that 22.1% of patients receiving pre-arrival care, including bystander CPR, survived, compared with 7.8% of patients that did not.

## **How AEDs Can Help**

Bystanders can further help a cardiac arrest victim's chances of survival by also using an AED. Brady helped spearhead an effort to install more than 300 AEDs across Grounds, helping UVA earn a HEARTSafe Campus award from the National Collegiate EMS Foundation twice in recent years.

Multiple research papers have shown the benefits of having AEDs accessible for use by members of the public. For example, a 2018 study of more than 50,000 out-of-hospital cardiac arrests found that 66.5% of patients who were treated with an AED by a bystander survived, compared with 43% of patients who were shocked only after fire rescue crews arrived. This highlights the benefits of quick action from bystanders. As Brady's article notes, "for every minute that a person with out-of-hospital cardiac arrest goes without CPR and defibrillation, the chance of survival decreases by 7 to 10%."

## **The Need for Education**

One major barrier to bystanders providing lifesaving care for cardiac arrest victims is a lack of training, especially in how to perform CPR. Brady's article points to a study that estimates only 2.4% of Americans receive CPR training each year.

One potential way to boost the number of people getting trained, Brady and his co-authors write, is to offer shorter training sessions. Potential

options include 4-minute training sessions at CPR kiosks in public locations, as well as video training sessions ranging from 60 seconds to 8 minutes.

Another option to boost bystander care is one Brady hopes will soon be available in Central Virginia: a smartphone app that notifies volunteer users of a cardiac arrest occurring near them in a public location. This would enable people close to the scene to start CPR and use an AED, if available, before emergency medical providers arrive. Albemarle County Fire Rescue, Charlottesville Fire Department, the Charlottesville-Albemarle-UVA Emergency Communications Center and UVA Health are working together to launch this tool.

"The next major advance in cardiac [arrest](#) management is not a new hospital-based treatment nor fire-rescue intervention. It is this pre-arrival care performed by a bystander, before fire-rescue personnel arrive and long before a patient is transported to the hospital. This care can allow victims of [cardiac arrest](#) to return to their lives and families," Brady said.

**More information:** William J. Brady et al, Lay Responder Care for an Adult with Out-of-Hospital Cardiac Arrest, *New England Journal of Medicine* (2019). [DOI: 10.1056/NEJMra1802529](https://doi.org/10.1056/NEJMra1802529)

Provided by University of Virginia

Citation: Bystanders can help more cardiac arrest victims survive (2019, December 5) retrieved 9 April 2024 from <https://medicalxpress.com/news/2019-12-bystanders-cardiac-victims-survive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---