

In some children with autism, 'social' and 'visual' neural circuits don't quite connect

December 17 2019



Credit: CC0 Public Domain

Among the first and most-documented symptoms of autism spectrum disorder (ASD) is a child's aversion to interaction with others. Specifically, they appear uninterested in social activities and stimuli that



would normally attract a young child's attention, such as watching other children play, sing or dance.

In a new study, published online December 17, 2019 in the journal eLife, researchers at University of California San Diego School of Medicine combined a novel vision tracking program with brain imaging to find that ASD toddlers who ignore social stimuli and prefer to look at moving, colorful geometric images, had more severe social symptoms and lower levels of brain activity connecting social and visual attention brain networks.

"This indicates that in a subtype of ASD toddlers with a preference for geometric images rather than pictures of children—about 20 percent of toddlers with autism—there is a disconnection between visual and social brain networks. In these ASD toddlers, colorful moving shapes, rather than fun social-emotional stimuli, control <u>neural activity</u>, attention and learning," said senior and corresponding author Karen Pierce, Ph.D., professor of neurosciences and co-director of the UC San Diego Autism Center of Excellence with co-author Eric Courchesne, Ph.D., also a professor of neurosciences.

The daily disconnection of social and visual neural networks results in attention, experience and learning to be directed towards low-level, but visually salient stimuli like colorful spinning shapes, said Pierce; this may be a causative factor in the symptoms and observed social impairment in some ASD toddlers. But the findings, she added, may also provide a new avenue for diagnosing and treating ASD in toddlers and young children.

"Currently, when a child receives a diagnosis of autism, he or she is usually referred for a fairly generic treatment based on principles of Applied Behavior Analysis. In the future, following a diagnosis, children might receive more in-depth biological evaluations that provide



information about their eye gaze and brain <u>network</u> activation patterns, which could point to more specific treatments," she said.

"It may also be possible that brain imaging and eye tracking could be used together to determine the efficacy of treatment if a second biological evaluation is conducted at some point following a period of treatment."

Pierce and colleagues have been investigating the potential of <u>eye gaze</u> technology in diagnosing and treating ASD for several years. In 2010, for example, they reported that, in a simple one minute eye tracking test, infants as young as 14 months who preferred movies of geometric shapes more often than movies of children dancing and doing yoga were subsequently diagnosed as ASD using a longer, gold-standard behavioral diagnostic test (the "ADOS"). Conversely, typically developing infants and toddlers preferred watching the "social" images.

In this new study, the researchers combined eye tracking (in which a camera monitors and documents where and what a child is looking at on a screen) with functional magnetic resonance imaging (fMRI) data detailing interconnectivity between different brain circuits.

"Basic neuroscience has found that the human brain has many so-called 'resting state' networks, each involved in different sensory, attention, cognitive and social functions. These networks are active even when we are not engaged in any explicit task, even during natural sleep," said Courchesne.

"One social network, the Default Mode Network or DMN, is highly active when we are thinking about ourselves and others. It is thought that abnormalities in the DMN may be central to why individuals with autism have social difficulties. Since experience-dependent mechanisms, such as what someone looks at, drives brain development, understanding what



someone visually attends—social or non-social stimuli—can provide invaluable information."

Examining the combined data for both ASD toddlers and non-ASD comparison groups, the researchers found less-than-typical neural interaction (hypoconnectivity) between social brain circuitry like the DMN and visual and attention networks in ASD children. The greater the hypoconnectivity, the more severe the social-communication difficulties, particularly in toddlers with geometric-preference ASD.

Pierce said the findings add new information and detail to the still largely mysterious and complex portrait of ASD. "But by combining clinical phenotype information, such as scores on tests of social competency, with brain imaging and eye tracking as we do here, we are developing more accurate, early approaches to diagnosing ASD and identifying brain-eye tracking subtypes. We will soon begin pilot studies to develop targeted treatments for this subtype."

More information: Michael V Lombardo et al, Default mode-visual network hypoconnectivity in an autism subtype with pronounced social visual engagement difficulties, *eLife* (2019). <u>DOI: 10.7554/eLife.47427</u>

Provided by University of California - San Diego

Citation: In some children with autism, 'social' and 'visual' neural circuits don't quite connect (2019, December 17) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2019-12-children-autism-social-visual-neural.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.