


Early regional anesthesia effective for extremity injury pain

December 17 2019



(HealthDay)—Regional anesthesia may aid pain management if used

soon after traumatic injury of the extremities, according to a study published in the December issue of *Regional Anesthesia & Pain Medicine*.

Rollin M. Gallagher, M.D., M.P.H., from the Corporal Michael J. Crescenz Veterans Affairs Medical Center in Philadelphia, and colleagues examined the use of [regional anesthesia](#) on combat-related extremity injury outcomes. The [analysis](#) included 358 American military personnel injured in Operation Enduring Freedom and Operation Iraqi Freedom (2007 to 2013) who were treated at two military treatment facilities and followed for two years after the injury.

The researchers found that receiving early regional anesthesia was associated with improved average pain during the first six months after injury when adjusting for injury severity and length of stay at the primary treatment facility. This benefit was sustained up to 24 months after injury. While individuals receiving early regional anesthesia reported greater pain relief, improved neuropathic pain intensity, and higher satisfaction with pain outcomes, these mean scores did not significantly differ between groups at 24 months.

"Findings indicate that when administered soon after traumatic injury, [regional anesthesia] is a valuable [pain management](#) intervention," the authors write.

More information: [Abstract/Full Text](#)

Copyright © 2019 [HealthDay](#). All rights reserved.

Citation: Early regional anesthesia effective for extremity injury pain (2019, December 17) retrieved 23 April 2024 from <https://medicalxpress.com/news/2019-12-early-regional-anesthesia-effective-extremity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.