


Research sheds important light on the metastasis of neuroblastoma

December 11 2019, by Paul Wiggins



Credit: Children's Cancer Institute Australia

Neuroblastoma can grow and spread in the body very aggressively, making it very difficult to treat. New research has shed important light on the factors at play in the metastasis of neuroblastoma, which could potentially lead to new treatment strategies.

The findings, recently published in the *British Journal of Cancer*, were discovered by Professor Maria Kavallaris and her team, through [scientific research](#) funded by The Kids Cancer Project.

The aggressive nature of neuroblastoma

Neuroblastoma is the most common solid tumour found in young children and is renowned for its aggressive nature—its ability to quickly grow and spread (metastasise) throughout the body. Sadly, children with [neuroblastoma](#) that has metastasised have a less than 50% chance of survival. There is an urgent need to better understand metastasis, so we might learn how to improve the treatment of metastatic neuroblastoma.

Understanding metastasis

Cancer spreads when [cells](#) become detached from the original tumour (the primary tumour), move to the surrounding tissue (migration), then get into the blood or lymphatic vessels where they can end up in different parts of the body (invasion). Metastasis is a complex process that involves many different steps and molecules in the body. In the study just published, Professor Kavallaris and her team have shown that a protein called stathmin may be a very important player in this process.

The role of stathmin

Stathmin is known to be overexpressed in neuroblastoma cells; that is, it occurs at higher levels in neuroblastoma cells than healthy cells. In earlier research, the team found that stathmin appears to help control the migration and invasion of neuroblastoma cells. In the new study, they tried to work out exactly how it does this, and how the process might be exploited to provide a therapeutic advantage. After all, if we can understand the factors that control metastasis, we may be able to inhibit these factors.

How stathmin does its work... and the significance

When they investigated stathmin's role in metastasis, the team found that it has a direct effect on a number of genes and other molecules called 'micro RNAs' (miRNAs) that are directly involved in the migration and invasion of neuroblastoma cells. In particular, stathmin appears to affect a protein called PTPN14, which has recently been identified as a tumour-suppressor.

"Our results showed that stathmin can regulate PTPN14 expression, which in turn affects neuroblastoma cell migration and invasion," explains Professor Kavallaris.

"This study has highlighted that both stathmin and PTPN14 appear to be important factors in metastasis of neuroblastoma. This raises the exciting possibility that targeting one or both of these molecules could be a potential new treatment strategy against this devastating disease."

"A substantial part of the initial project was supported by funding from The Kids' Cancer Project, which was invaluable in allowing this work to progress," said Professor Kavallaris.

More information: Sela T. Po'uha et al, Stathmin levels alter PTPN14 expression and impact neuroblastoma cell migration, *British Journal of*

Cancer (2019). [DOI: 10.1038/s41416-019-0669-1](https://doi.org/10.1038/s41416-019-0669-1)

Provided by Children's Cancer Institute Australia

Citation: Research sheds important light on the metastasis of neuroblastoma (2019, December 11) retrieved 25 April 2024 from <https://medicalxpress.com/news/2019-12-important-metastasis-neuroblastoma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.