

Report links recommended physical activity levels to lower risk of seven cancers

December 26 2019



Credit: CC0 Public Domain

A pooled analysis of nine prospective studies involving more than 750,000 adults finds that recommended amounts of leisure-time physical activity were linked to a lower risk for seven cancers, with several cancer types having a 'dose/response' relationship. The study was led by investigators at the National Cancer Institute, the American Cancer

Society, and the Harvard T.H. Chan School of Public Health and appears in the *Journal of Clinical Oncology*.

While it's long been known that [physical activity](#) is associated with a [lower risk](#) of several cancers, less clear has been the shape of the relationship and whether recommended amounts of physical activity are associated with lower risk. Updated guidelines for activity now state that people should aim for 2.5 to 5 hours/week of moderate-intensity activity or 1.25 to 2.5 hours/week of vigorous activity. Moderate-intensity activities are those that get you moving fast enough or strenuously enough to burn off three to six times as much energy per minute as sitting quietly (3 to 6 METs). Vigorous-intensity activities burn more than 6 METs.

For the current analysis, investigators pooled data from nine prospective cohorts with self-reported leisure-time physical activity and follow-up for cancer incidence, looking at the relationship between physical activity with incidence of 15 types of cancer.

They found engaging in recommended amounts of activity (7.5 to 15 MET hours/week) was associated with a statistically significant lower risk of seven of the 15 [cancer types](#) studied, with the reduction increasing with more MET hours. Physical activity was associated with a lower risk of colon cancer in men (8% for 7.5 MET hours/week; 14% for 15 MET hours/week), female breast cancer (6%-10%), endometrial cancer (10%-18%), kidney cancer (11%-17%), myeloma (14%-19%), liver cancer (18%-27%), and non-Hodgkin lymphoma (11%-18% in women). The dose response was linear in shape for half of the associations and nonlinear for the others.

The analysis had some limitations: Even with 750,000 participants, patient numbers were limited for some cancers; participants were primarily white; there was a limited number of cohorts with detailed

physical activity measures; and the authors relied on self-reported physical activity.

The authors conclude: "These findings provide direct quantitative support for the levels of activity recommended for cancer prevention and provide actionable evidence for ongoing and future cancer prevention efforts."

"Physical activity guidelines have largely been based on their impact on [chronic diseases](#) like [cardiovascular disease](#) and diabetes," said Alpa Patel, Ph.D., senior scientific director of epidemiology research at the American Cancer Society. "These data provide strong support that these recommended levels are important to [cancer prevention](#), as well."

More information: Amount and Intensity of Leisure-Time Physical Activity and Lower Cancer Risk, Matthews et al. *Journal of Clinical Oncology* (2019). [DOI: 10.1200/JCO.19.02407](https://doi.org/10.1200/JCO.19.02407)

Provided by American Cancer Society

Citation: Report links recommended physical activity levels to lower risk of seven cancers (2019, December 26) retrieved 23 April 2024 from <https://medicalxpress.com/news/2019-12-links-physical-cancers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.