


A molecular map of the brain's decision-making area

December 24 2019



From left: Konstantinos Meletis, Antje Märtin, Daniela Calvigioni and Rania Tzortzi, researchers at the Department of Neuroscience at Karolinska Institutet. Credit: Juan Perez Fernandez

Researchers at Karolinska Institutet have come one step closer toward understanding how the part of our brain that is central for decision-making and the development of addiction is organized on a molecular level. In mouse models and with methods used for mapping cell types and brain tissue, the researchers were able to visualize the organization of different opioid-islands in striatum. Their spatiomolecular map, published in the journal *Cell Reports*, may further our understanding of the brain's reward-system.

Striatum is the inner part of the brain that among other things regulates rewards, motivation, impulses and motor function. It is considered central to decision-making and the development of various addictions.

In this study, the researchers created a molecular 3-D-map of the nerve cells targeted by opioids, such as morphine and heroin, and showed how they are organized in [striatum](#). It is an important step toward understanding how the brain's network governing motivation and drug addiction is organized. In the study, the researchers described a spatiomolecular code that can be used to divide striatum into different subregions.

"Our map forms the basis for a new understanding of the brain's probably most important network for decision-making," says Konstantinos Meletis, associate professor at the Department of Neuroscience at Karolinska Institutet and the study's main author. "It may contribute to an increased understanding of both normal reward processes and the effects of various addictive substances on this network."

To find this molecular code, the researchers used single-nucleus RNA sequencing, a method to study small differences in individual cells, and mapping of the striatal gene expression. The results provide the first demonstration of molecular codes that divide the striatum into three

main levels of classification: a spatial, a patch-matrix and a cell-type specific organization.

"With this [new knowledge](#) we may now begin to analyze the function of different types of nerve [cells](#) in different molecularly defined areas," says Meletis. "This is the first step in directly defining the networks' role in controlling decision-making and addiction with the help of optogenetics."

This new knowledge may also form the basis for the development of new treatments based on a mechanistic understanding of the brain's network, according to the researchers.

More information: "A spatiomolecular map of the striatum," Antje Märtin, Daniela Calvigioni, Ourania Tzortzi, Janos Fuzik, Emil Wärnberg, Konstantinos Meletis, *Cell Reports*, online Dec. 24, 2019, [DOI: 10.1016/j.celrep.2019.11.096](https://doi.org/10.1016/j.celrep.2019.11.096)

Provided by Karolinska Institutet

Citation: A molecular map of the brain's decision-making area (2019, December 24) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2019-12-molecular-brain-decision-making-area.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--