

Researchers identify neurons responsible for rapid eye movements/REM during sleep

December 20 2019



Credit: CC0 Public Domain

Rapid Eye Movement—not only the name of a successful American rock band, but also and not least a characteristic eye movement in paradoxical sleep, the stage with high dream activity. This sleep phase



has a peculiarity: Although the muscle tone of the sleeping person is completely relaxed, the eyes suddenly move back and forth. The name "paradoxical sleep" is well deserved. Characteristic of these are signs of deep sleep (muscle atony) in connection with brain activity, which is very similar to those in the waking state and its eye movements. This sleep phase was discovered in the 1950s by French and American researchers and consequently called rapid eye movement sleep (REM sleep), i.e. sleep with rapid eye movements. Why can this strange phenomenon be useful? For 70 years, scientists have been dreaming of getting to the bottom of the mystery. Thanks to the productive cooperation between the universities of Bern and Fribourg, this dream could now come true.

Butterfly wings arranged neurons

For several years, the team led by Franck Girard and Marco Celio at the University of Fribourg has studied <u>neurons</u> under the microscope, that occur in the <u>brain stem</u> and form a structure that is reminiscent of butterfly wings, which is why the structure is called Nucleus papilio. "These neurons are associated with multiple nerve centers, especially those responsible for eye <u>movement</u>, and those involved in sleep control," explains Franck Girard. "Therefore, we asked ourselves the following question: may the nucleus papilio neurons play a role in the control of eye movements during sleep?"

Stronger together

To test this hypothesis, the Fribourg researchers turned to the research group headed by Dr. C. Gutiérrez Herrera and Prof. A. Adamantidis at the Department of Neurology at the Inselspital, University Hospital Bern, and Department for BioMedical Research of the University of Bern, who are investigating sleep in mice. "To our surprise, we found



that these neurons are particularly active in the phase of paradoxical sleep," reports Dr. Carolina Gutierrez. The researchers from Bern gathered the loop around the nucleus papilio neurons even more closely and were able to demonstrate with the help of optogenetic methods (combined optical and genetic techniques) that their artificial activation causes <u>rapid eye movement</u>, especially during this <u>sleep phase</u>. Conversely, the inhibition or elimination of these same neurons blocks the movement of the eyes.

After the "how" the "why"

Now that it is clear that the nucleus papilio neurons play an important role in eye movement during REM sleep, it is important to find out what function this phenomenon has. Is it due to the visual experience of dreams? Does it matter in preserving memories? "Now that we are able to specifically activate the nucleus papilio 'on demand' in mice by optogenetic methods, we may be able to find answers to these questions," says Antoine Adamantidis. The next step, however, will be to confirm the activation of nucleus papilio neurons during REM sleep in humans. The researchers have not yet found the key to their dreams, but they've come a long way.

A better understanding of the neural circuits involved in paradoxical sleep is therefore a prerequisite for understanding for instance how these neurons are prone to degenerative changes in diseases such as Parkinson's.

More information: C. Gutierrez Herrera et al, Neurons in the Nucleus papilio contribute to the control of eye movements during REM sleep, *Nature Communications* (2019). DOI: 10.1038/s41467-019-13217-y



Provided by University of Bern

Citation: Researchers identify neurons responsible for rapid eye movements/REM during sleep (2019, December 20) retrieved 15 May 2024 from https://medicalxpress.com/news/2019-12-neurons-responsible-rapid-eye-movementsrem.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.