

Permanent hair dye and straighteners may increase breast cancer risk

December 4 2019



The study found that women who use permanent hair dye and chemical hair straighteners have a higher risk of developing breast cancer than women who don't use these products. Credit: NIEHS

Scientists at the National Institutes of Health found that women who use

permanent hair dye and chemical hair straighteners have a higher risk of developing breast cancer than women who don't use these products. The study published online Dec. 4 in the *International Journal of Cancer* and suggests that breast cancer risk increased with more frequent use of these chemical hair products.

Using data from 46,709 women in the Sister Study, researchers at the National Institute of Environmental Health Sciences (NIEHS), part of NIH, found that women who regularly used permanent hair dye in the year prior to enrolling in the study were 9% more likely than women who didn't use hair dye to develop breast [cancer](#). Among African American women, using permanent dyes every five to eight weeks or more was associated with a 60% increased risk of breast cancer as compared with an 8% increased risk for white women. The research team found little to no increase in [breast cancer risk](#) for semi-permanent or temporary dye use.

"Researchers have been studying the possible link between hair dye and cancer for a long time, but results have been inconsistent," said corresponding author Alexandra White, Ph.D., head of the NIEHS Environment and Cancer Epidemiology Group. "In our study, we see a higher breast cancer risk associated with hair dye use, and the effect is stronger in African American women, particularly those who are frequent users."

An intriguing finding was the association between the use of chemical hair straighteners and breast cancer. Dr. White and colleagues found that women who used hair straighteners at least every five to eight weeks were about 30% more likely to develop breast cancer. While the association between straightener use and breast cancer was similar in African American and [white women](#), straightener use was much more common among African American women.

Co-author Dale Sandler, Ph.D., chief of the NIEHS Epidemiology Branch, cautioned that although there is some prior evidence to support the association with chemical straighteners, these results need to be replicated in other studies.

When asked if women should stop dyeing or straightening their hair, Sandler said, "We are exposed to many things that could potentially contribute to breast cancer, and it is unlikely that any single factor explains a woman's risk. While it is too early to make a firm recommendation, avoiding these chemicals might be one more thing [women](#) can do to reduce their risk of [breast cancer](#)."

More information: Carolyn E. Eberle et al, Hair dye and chemical straightener use and breast cancer risk in a large US population of black and white women, *International Journal of Cancer* (2019). [DOI: 10.1002/ijc.32738](#)

Provided by National Institutes of Health

Citation: Permanent hair dye and straighteners may increase breast cancer risk (2019, December 4) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-12-permanent-hair-dye-straighteners-breast.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--