

# Personalized medicine for atrial fibrillation

December 16 2019

---



From left to right: José Manuel Alfonso Almazán, Jorge García Quintanilla, Manuel Marina Breyse, José María Lillo Castellano, Julián Pérez Villacastín, David Filgueiras Rama, José Jalife y Nicasio Pérez Castellano. Credit: CNIC

Patients with atrial fibrillation, the most frequent cardiac arrhythmia, are closer to accessing personalized medicine. This is the claim of a new study led by Dr. David Filgueiras, of the Centro Nacional de

Investigaciones Cardiovasculares (CNIC), the Hospital Clínico San Carlos de Madrid, and the Centro de Investigación en Red de Enfermedades Cardiovasculares (CIBERCV). The study, results show that it is possible to monitor and predict individual progression of atrial fibrillation from cardiac electrical signals obtained from implantable devices (pacemakers or defibrillators). This project has been supported by the Fundación Interhospitalaria para la Investigación Cardiovascular (Fundación FIC).

The study, published in the latest edition of *Europace*, involved the participation of 51 Spanish hospitals and the Fundación Interhospitalaria para la Investigación Cardiovascular. The results show that cardiac electrical signals from patients fitted with pacemakers or [implantable cardioverter-defibrillators](#) can be used to monitor and predict the progression of the arrhythmia in a personalized and specific manner.

This is achieved with standard data transmission technology installed in implantable devices such as pacemakers. This technology can be used to monitor cardiac electrical activity during episodes of atrial [fibrillation](#), thus establishing disease status and the rate of progression.

Atrial fibrillation is the most frequent chronic sustained arrhythmia, affecting more than 30 million people worldwide and an estimated 600,000 or more people in Spain. One in every three people will develop the condition in their lifetime. The risk is higher in people with certain diseases, such as diabetes or hypertension. Risk is also increased by lifestyle habits such as excessive consumption of alcohol or other substances.

Atrial fibrillation, explained Dr. Filgueiras, tends to begin with episodes of short duration that resolve themselves. As the condition progresses, episodes become more prolonged until they become persistent. At this stage, episodes no longer terminate spontaneously, and the heart

struggles to recover a healthy, stable rhythm, even after medication or surgery.

In recent years, several risk scales have been developed to estimate individual risk of developing atrial fibrillation over a 10-year period. However, according to Dr. Filgueiras, "once atrial fibrillation develops, its progression varies significantly between patients; in some, a pattern of repeated short episodes is maintained for years, whereas in others it progresses rapidly to longer episodes associated with atrial dilatation and major difficulties in controlling heart rhythm."

There is thus a need for methods to individually monitor and predict the progression of atrial fibrillation. This would allow, after the initiation of atrial fibrillation episodes, "optimal timing of the most appropriate medical treatment, avoiding delays until the arrhythmia has progressed to a state that is harder to control," explained CNIC researcher Dr. José Jalife, an author on the study.

According to Dr. Filgueiras, "this technology opens up enormous possibilities in [personalized medicine](#) for atrial fibrillation patients because it allows us to determine the progression rate of the arrhythmia in each individual and to optimize the timing of medical intervention with current treatment options."

In addition, Dr. Julián Villacastín, director of the Instituto Cardiovascular del Hospital Clínico San Carlos and an author on the study, this new approach to atrial fibrillation diagnosis "will allow us to monitor the influence of different interventions on disease progression."

The study findings were obtained by independent big-data screening of databases held at leading national hospitals in Spain together with interdisciplinary collaboration between physicians and engineers, including first author Dr. José María Lillo of the CNIC and the

Fundación Interhospitalaria para la Investigación Cardiovascular. Dr. Lillo explained that "tools based on signals from pacemakers or defibrillators could in the near future be adapted to exploit signals from other devices such as smart watches or even holter devices for long-duration ECG recording." This has the potential to expand this type of personalized medicine to a much larger population of patients with conditions other than [atrial fibrillation](#).

**More information:** José María Lillo-Castellano et al, Personalized monitoring of electrical remodelling during atrial fibrillation progression via remote transmissions from implantable devices, *EP Europace* (2019). [DOI: 10.1093/europace/euz331](https://doi.org/10.1093/europace/euz331)

Provided by Centro Nacional de Investigaciones Cardiovasculares Carlos III (F.S.P.)

Citation: Personalized medicine for atrial fibrillation (2019, December 16) retrieved 3 May 2024 from <https://medicalxpress.com/news/2019-12-personalized-medicine-atrial-fibrillation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--