

Three quarters of teens who vape report using nicotine, marijuana, or multiple substances

December 16 2019



Credit: CC0 Public Domain

More teens who vape are using addictive or mind-altering substances than previously believed, according to a new study in the *American Journal of Preventive Medicine*. The data paint a different picture than

previous research because of the significantly higher proportion (75 percent) of teens who vape using nicotine, marijuana, or multiple substances and not just flavoring. These findings add to growing public health concerns about youth vaping.

"Our study provided a more nuanced view of youth e-cigarette use behaviors than earlier studies. We found that youth were more likely to report [vaping nicotine](#) and marijuana than 'just flavoring' only, and that cigarette smoking intensity was associated with an increasing proportion of students reporting vaping nicotine only," explained co-investigator Hongying Dai, Ph.D., Associate Professor, College of Public Health, University of Nebraska Medical Center, Omaha, NE, USA.

This study examined patterns of youth vaping nicotine, marijuana, and just flavoring in the past 30 days by analyzing data from the 2017 Monitoring the Future (MTF) cross-sectional study. Of the 14,560 teens participating in that study, 12 percent reported vaping within the prior 30 days, with 7.4 percent using nicotine and 3.6 percent, marijuana. Of that group, only 24.9 percent reported vaping just flavoring only, while a majority (75.1 percent) reported vaping nicotine, marijuana, or multiple substances.

Current cigarette smoking intensity was associated with an [increased risk](#) of reporting vaping all three substances. Compared with 8th graders, more 10th and 12th graders reported vaping nicotine, [marijuana](#), and just flavoring during the study period. Female students were also less likely to report vaping these three substances than male students. Fewer non-Hispanic blacks reported vaping nicotine and just flavoring than non-Hispanic whites. Hispanics were also less likely to report vaping nicotine.

The prevalence of e-cigarette use among US youth increased dramatically during 2017-2019, partly due to the rising popularity of

products with nicotine salt and pod-based products like JUUL, and a large number of flavors appealing to adolescents. The nationwide increases in use led the US Surgeon General to issue an advisory about the epidemic in 2018, but much more needs to be done to reverse the upward trend. Due to the recent spate of vaping-related lung injuries, calls for restrictions on flavored vaping products and e-cigarette use have become more urgent. Identifying [substances](#) vaped by youth is critical to formulating, implementing, and evaluating population-wide strategies and interventions to curb youth use of these products.

"Continuous surveillance of youth behaviors and strategies and interventions to reduce [youth](#) e-cigarette use are needed. The truth is that no form of tobacco is safe," added co-investigator Mohammad Siahpush, Ph.D., College of Public Health, University of Nebraska Medical Center, Omaha, NE, USA.

More information: Hongying Dai et al, Use of E-Cigarettes for Nicotine, Marijuana, and Just Flavoring Among U.S. Youth, *American Journal of Preventive Medicine* (2019). [DOI: 10.1016/j.amepre.2019.09.006](#)

Provided by Elsevier

Citation: Three quarters of teens who vape report using nicotine, marijuana, or multiple substances (2019, December 16) retrieved 24 May 2024 from <https://medicalxpress.com/news/2019-12-quarters-teens-vape-nicotine-marijuana.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
