

Reducing the side-effects of prostate hormone therapy with exercise

December 9 2019

A prescription of short-term exercise for patients with advanced prostate cancer could help to reduce the side-effects of hormone therapy, according to new research from the University of East Anglia.

Researchers from the Norfolk and Norwich University Hospital (NNUH) and UEA led a trial which involved patients who were due to start [androgen deprivation therapy](#) (ADT).

Fifty patients took part in the research study, with half of the participants taking part in two supervised [exercise](#) sessions a week for three months at specialist exercise science facilities at UEA.

The trial aimed to reduce the adverse side-effects of hormone therapy such as [weight gain](#) and an increased risk of heart problems and assessed participants' health three months after their exercise programme.

The findings, which have been published in the *British Journal of Urology International (BJUI)*, showed that the three month programme of aerobic and resistance training intervention prevented adverse changes in cardiopulmonary fitness and fatigue for those taking part in the trial.

Prof John Saxton, from UEA's Norwich Medical School, said:
"Structured exercise programmes have much to offer people living with common cancers. Hormone treatments for prostate cancer are known to have undesirable side-effects, which increase a man's susceptibility to [cardiovascular disease](#).

"This research shows that some of the harmful side-effects of hormone therapy are reduced in men who begin to exercise regularly around the same time that these drugs are prescribed.

"Our findings have important implications for the quality of prostate cancer survival," he added.

Wilphard Ndjavera, Clinical Fellow for Urology at NNUH, said: "All the patients really enjoyed the sessions and the results have suggested that exercise does work to reduce these harmful side effects of treatment. It is one of only two studies in the world to look at this."

"After the supervised exercise was withdrawn, differences in cardiopulmonary fitness and fatigue were not sustained, but the exercise group showed higher quality of life and reduced cardiovascular risk compared to the [control group](#)."

"These findings have important implications for clinicians concerned with the management of ADT-related side-effects."

"Exercise-induced attenuation of treatment side-effects in patients with newly diagnosed [prostate cancer](#) beginning androgen-deprivation therapy: a randomised controlled trial" is published in the *British Journal of Urology International (BJUI)*.

More information: Wilphard Ndjavera et al. Exercise-induced attenuation of treatment side-effects in patients with newly diagnosed prostate cancer beginning androgen-deprivation therapy: a randomised controlled trial, *BJU International* (2019). [DOI: 10.1111/bju.14922](https://doi.org/10.1111/bju.14922)

Provided by University of East Anglia

Citation: Reducing the side-effects of prostate hormone therapy with exercise (2019, December 9) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2019-12-side-effects-prostate-hormone-therapy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.