

New treatment strategy may thwart deadly brain tumors

December 23 2019



Killer T cells surround a cancer cell. Credit: NIH

Immune checkpoint inhibitors are important medications that boost the immune system's response against certain cancers; however, they tend to be ineffective against glioblastoma, the most deadly primary brain tumor

in adults. New research in mice led by investigators at Massachusetts General Hospital (MGH) and the University of Florida reveals a promising strategy that makes glioblastoma susceptible to these medications. The findings, which are published in the *Proceedings of the National Academy of Sciences*, indicate that such combination therapy should be tested in clinical trials of patients with glioblastoma, for whom there is no known cure.

Part of the reason glioblastoma does not respond well to [immune checkpoint inhibitors](#) and other immunotherapies is because cells called myeloid-derived suppressor cells (MDSCs) infiltrate the region surrounding glioblastoma tumors, where they contribute to immunosuppression, tumor progression, and treatment resistance. Thus, targeting these cells may augment immunotherapy and improve responses to treatment in affected patients.

A [collaborative effort](#) co-led by Jeffrey K. Harrison, Ph.D., of the Department of Pharmacology and Therapeutics at the University of Florida, and Rakesh K. Jain, Ph.D., of the Department of Radiation Oncology at MGH and Harvard Medical School, set out to test this strategy. Using two mouse models of glioblastoma, the team targeted receptors—called chemokine receptors—that are important for allowing MDSCs to infiltrate into the region surrounding glioblastoma tumors. In mice that were bred to lack chemokine receptor 2 (CCR2) and to develop glioblastoma, MDSCs could not carry out such infiltration. Treating these mice with an immune checkpoint inhibitor stimulated a strong anti-cancer immune response and prolonged the animals' survival. In [mice](#) with normal CCR2, treatment with a molecule that blocks CCR2 had similar effects.

"The CCR2 antagonist used in this study—called CCX872— has passed phase Ib safety trials in patients with pancreatic tumors, and [clinical trials](#) are ongoing to investigate the use of CCR2 inhibitors in several

cancers," said Jain. "Thus, the results of this study support targeting CCR2-expressing MDSCs as a means to enhance immunotherapies, and warrant investigation of this combination therapy in clinical trials for patients with glioblastoma."

More information: Joseph A. Flores-Toro et al., "CCR2 inhibition reduces tumor myeloid cells and unmasks a checkpoint inhibitor effect to slow progression of resistant murine gliomas," *PNAS* (2019).

www.pnas.org/cgi/doi/10.1073/pnas.1910856117

Provided by Massachusetts General Hospital

Citation: New treatment strategy may thwart deadly brain tumors (2019, December 23) retrieved 18 April 2024 from

<https://medicalxpress.com/news/2019-12-treatment-strategy-thwart-deadly-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.