

Low cardiac troponin levels cannot safely rule out ischemia

January 7 2020



(HealthDay)—Even very low concentrations of high-sensitivity cardiac

troponin I (hs-cTnI) cannot safely rule out inducible myocardial ischemia in patients with symptomatic coronary artery disease (CAD), according to a study published online Jan. 7 in the *Annals of Internal Medicine*.

Joan Walter, M.D., Ph.D., from University Hospital Basel in Switzerland, and colleagues measured circulating hs-cTn concentrations among 1,896 consecutive patients with CAD referred with symptoms possibly related to inducible myocardial ischemia. Providers were blinded to results from tests used to adjudicate the presence of inducible myocardial ischemia, including [myocardial perfusion](#) imaging with single-photon emission computed tomography and, when available, [coronary angiography](#) and fractional flow reserve measurements.

The researchers found that 46 percent of patients had inducible myocardial ischemia. A cutoff value of 2.5 ng/L for hs-cTnI provided a negative predictive value (NPV) of 70 percent and a sensitivity of 90 percent for exclusion of inducible myocardial ischemia. There was no hs-cTnI cutoff that reached both predefined performance characteristic targets. With alternative assays (hs-cTnI or hs-cTnT), there was no cutoff that achieved the target performance: hs-cTnT concentrations

Citation: Low cardiac troponin levels cannot safely rule out ischemia (2020, January 7) retrieved 10 April 2024 from <https://medicalxpress.com/news/2020-01-cardiac-troponin-safely-ischemia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
