

Cyberbullying linked to increased depression and PTSD

January 22 2020



Philip D. Harvey, Ph.D, professor of psychiatry and behavioral sciences at the University of Miami Miller School of Medicine, co-author of "Cyberbullying and Its Relationship to Current Symptoms and History of Early Life Trauma." Credit: University of Miami Health System

Cyberbullying had the impact of amplifying symptoms of depression and

post-traumatic stress disorder in young people who were inpatients at an adolescent psychiatric hospital, according to a new study published in the *Journal of Clinical Psychiatry*. The study addressed both the prevalence and factors related to cyberbullying in adolescent inpatients.

"Even against a backdrop of emotional challenges in the kids we studied, we noted cyberbullying had an adverse impact. It's real and should be assessed," said Philip D. Harvey, Ph.D., professor of psychiatry and behavioral sciences at the University of Miami Miller School of Medicine, who co-authored the paper "Cyberbullying and Its Relationship to Current Symptoms and History of Early Life Trauma."

He says children with a history of being abused were found to be more likely to be cyberbullied, suggesting that assessments for [childhood trauma](#) should also include assessments for cyberbullying. Likewise, children who report being cyberbullied should be assessed for a history of childhood trauma.

"Cyberbullying is possibly more pernicious than other forms of bullying because of its reach," Dr. Harvey says. "The bullying can be viral and persistent. To really be bullying, it has to be personal—a directly negative comment attempting to make the person feel bad."

The study helped to confirm other facts about cyberbullying:

- Being online regularly or the amount of time spent on social media weren't determining factors in who was cyberbullied.
- Cyberbullying cuts across all economic classes and ethnic backgrounds.
- Adolescents who have been bullied in the past had a higher risk of being bullied again.

Studying Cyberbullying's Impact on an Inpatient

Psychiatric Population

The study of 50 adolescent psychiatric inpatients ages 13 to 17 examined the prevalence of cyberbullying and related it to social media usage, current levels of symptoms and histories of adverse early life experience.

Conducted from September 2016 to April 2017 at a suburban psychiatric hospital in Westchester County, New York, the study asked participants to complete two childhood trauma questionnaires and a cyberbullying questionnaire.

Twenty percent of participants reported that they had been cyberbullied within the last two months before their admission. Half of the participants were bullied by text messages and half on Facebook. Transmitted pictures or videos, Instagram, instant messages and chat rooms were other cyberbullying vehicles.

Those who had been bullied had significantly higher severity of post-[traumatic stress disorder](#) (PTSD), depression, anger, and fantasy dissociation than those who were not bullied.

Links to Childhood Trauma

Participants who reported being cyberbullied also reported significantly higher levels of lifetime emotional abuse on the study's Childhood Trauma Questionnaire than those who were not bullied. These same [young people](#) did not report a significantly higher level of other types of trauma ([physical abuse](#), sexual abuse, emotional neglect or physical neglect).

Further studies are needed to establish whether there may be some unique consequence of childhood [emotional abuse](#) that makes troubled

teens more likely to experience or report [cyberbullying](#).

Conclusions

While all of the participants in this study were psychiatric inpatients, those who had been bullied had significantly higher scores on PTSD, depression, anger, and dissociation scales than those who were not bullied. Dr. Harvey says this finding is consistent with past research.

Dr. Harvey encourages psychologists, psychiatrists and other counselors to routinely ask young people if they were abused or traumatized when they were younger and whether they are being bullied now.

He says adding these questions to the clinical evaluation of adolescents may bring to light symptoms that may have otherwise been ignored. Additionally, factors that may be causing or contributing to those symptoms can be targeted for specific intervention.

Parents and adolescents can take action to discourage bullying, Dr. Harvey says. "It's not hard to block someone on the Internet, whether it's texting, Facebook, Twitter, or sending pictures. Ask why are people choosing you to bully? If it's something you're posting, assess that and make a change."

Dr. Harvey collaborated on the study with Samantha Block Saltz, M.D., also with the University of Miami Miller School of Medicine, and David L. Pogge, Ph.D., of Winds Hospital in Katonah, New York.

Provided by University of Miami Leonard M. Miller School of Medicine

Citation: Cyberbullying linked to increased depression and PTSD (2020, January 22) retrieved 23 April 2024 from

<https://medicalxpress.com/news/2020-01-cyberbullying-linked-depression-ptsd.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.