


Depression common during perimenopause but not regularly assessed by providers

January 22 2020



Credit: CC0 Public Domain

Almost 40% of women experience symptoms of depression during perimenopause, yet it often goes undetected and untreated because many healthcare providers aren't screening for it and aren't prepared with

treatment options. A new study analyzed screening practices by women's healthcare providers and their management of depression during perimenopause. Study results are published online in *Menopause*, the journal of The North American Menopause Society (NAMS).

The high number of women experiencing symptoms of [depression](#) has been linked to hormone changes, historical depressive episodes, [life events](#), and a genetic predisposition to depression. As more data emerges to support the presence of risk factors for depression during [perimenopause](#) and as the patient population continues to age, [healthcare providers](#) need to become better educated in order to recognize and manage depression.

Rates of routine [screening](#) and healthcare-provider beliefs and knowledge about symptoms of depression in perimenopause remained unstudied until now. The aim of this new study was to better understand clinical practice patterns of obstetrician-gynecologists regarding their screening practices and management of depression.

The survey was sent to 500 practicing obstetrician-gynecologists who were fellows of the American College of Obstetricians and Gynecologists and members of the Collaborative Ambulatory Research Network. It found that more than a third of respondents (34.1%) did not regularly screen perimenopausal women for depression and that more than half (55%) believed they were not equipped to manage the problem.

Higher-quality education about depression was associated with higher rates of screening. These findings suggest that improved training for obstetrician-gynecologists in the diagnosis, treatment, and management of depression, both in residency and later, could improve rates of screening.

Study results appear in the article "Obstetrician-gynecologists' screening

and management of depression during perimenopause."

"Given the prevalence of depressive symptoms in perimenopausal women, the recent publication of guidelines for the diagnosis and management of depression in this population, and the availability of safe and effective therapies, all healthcare providers should be screening their [female patients](#) for depression. Improving training for healthcare providers in the evaluation and treatment of depression is important to optimizing patient outcomes," says Dr. Stephanie Faubion, NAMS medical director.

More information: Greta B. Raglan et al, Obstetrician-gynecologists' screening and management of depression during perimenopause, *Menopause* (2020). [DOI: 10.1097/GME.0000000000001488](https://doi.org/10.1097/GME.0000000000001488)

Provided by The North American Menopause Society

Citation: Depression common during perimenopause but not regularly assessed by providers (2020, January 22) retrieved 27 April 2024 from <https://medicalxpress.com/news/2020-01-depression-common-perimenopause-regularly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.