


Groups publish statements on CT contrast use in patients with kidney disease

January 21 2020



Credit: CC0 Public Domain

The risk of administering modern intravenous iodinated contrast media in patients with reduced kidney function has been overstated, according

to new consensus statements from the American College of Radiology (ACR) and the National Kidney Foundation (NKF), published in the journal *Radiology*.

Intravenous iodinated [contrast media](#) are commonly used with computed tomography (CT) to evaluate disease and to determine treatment response. Although patients have benefited from their use, iodinated [contrast](#) media have been denied or delayed in patients with reduced [kidney function](#) due to the perceived risks of contrast-induced [acute kidney injury](#). This practice can hinder a timely and accurate diagnosis in these patients.

"The historical fears of [kidney injury](#) from contrast-enhanced CT have led to unmeasured harms related to diagnostic error and diagnostic delay," said lead author Matthew S. Davenport, M.D., associate professor of radiology and urology at the University of Michigan in Ann Arbor, Michigan. "Modern data clarify that this perceived risk has been overstated. Our intent is to provide multi-disciplinary guidance regarding the true risk to patients and how to apply a consideration of that risk to modern [clinical practice](#)."

These consensus statements were developed to improve and standardize the care of patients with impaired kidney function who may need to undergo exams that require intravenous iodinated contrast media to provide the clearest images and allow for the most informed diagnosis.

In clinical practice, many factors are used to determine whether intravenous contrast media should be administered. These include probability of an accurate diagnosis, alternative methods of diagnosis, risks of misdiagnosis, expectations about kidney function recovery, and risk of allergic reaction. Decisions are rarely based on a single consideration, such as risk of an adverse event specifically related to kidney impairment. Consequently, the authors advise that these

statements be considered in the context of the entire clinical scenario.

Importantly, the report outlines the key differences between contrast-induced acute kidney injury (CI-AKI) and contrast-associated acute kidney injury (CA-AKI). In CI-AKI, a causal relationship exists between contrast media and kidney injury, whereas in CA-AKI, a direct causal relationship has not been demonstrated. The authors suggest that studies that have not properly distinguished the two have contributed to the overstatement of risk.

"A primary explanation for the exaggerated perceived nephrotoxic risk of contrast-enhanced CT is nomenclature," Dr. Davenport said.

"'Contrast-induced' acute kidney injury implies a causal relationship. However, in many circumstances, the diagnosis of CI-AKI in clinical care and in research is made in a way that prevents causal attribution. Disentangling contrast-induced AKI (causal AKI) from contrast-associated AKI (correlated AKI) is a critical step forward in improving understanding of the true risk to patients."

The statements answer key questions and provide recommendations for use of intravenous contrast media in treating patients with varying degrees of impaired kidney function.

Although the true risk of CI-AKI remains unknown, the authors recommend intravenous normal saline for patients without contraindication, such as heart failure, who have acute kidney injury or an estimated glomerular filtration rate (eGFR) less than 30 mL/min per 1.73 m² who are not undergoing maintenance dialysis. In individual and unusual high-risk circumstances (patients with multiple comorbid risk factors), prophylaxis may be considered in patients with an eGFR of 30-44 mL/min per 1.73 m² at the discretion of the ordering clinician.

The presence of a solitary kidney should not independently influence

decision making regarding the risk of CI-AKI. Lowering of contrast media dose below a known diagnostic threshold should be avoided due to the risk of lowering diagnostic accuracy. Also, when feasible, medications that are toxic to the kidneys should be withheld by the referring clinician in patients at high risk. However, renal replacement therapy should not be initiated or altered solely based on contrast media administration.

The authors emphasize that prospective controlled data are needed in adult and pediatric populations to clarify the risk of CI-AKI.

More information: "Use of Intravenous Iodinated Contrast Media in Patients with Kidney Disease: Consensus Statements from the American College of Radiology and the National Kidney Foundation," *Radiology*, 2020.

Provided by Radiological Society of North America

Citation: Groups publish statements on CT contrast use in patients with kidney disease (2020, January 21) retrieved 6 May 2024 from <https://medicalxpress.com/news/2020-01-groups-publish-statements-ct-contrast.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
