

Possible key to beating colorectal cancer

January 29 2020



SRR-mediated dehydration of serine contributes to the pyruvate pool in colon cancer cells, maintains mitochondrial mass and leads to anti-apoptotic effect, resulting in enhancement of growth of colon cancer cells. Credit: Osaka University



Serine racemase (SRR) is a multifunctional enzyme that carries out several different reactions in human cells, including the conversion of Lserine into pyruvate. Though already well characterized, a team of researchers led by Osaka University has discovered an important new role for SRR in cancer metabolism, exposing the metabolic pathway as a viable target for novel anti-cancer therapies.

Researchers have long known that cancer cells display an altered metabolism that favors their growth, survival, and metastasis. Colorectal cancer is one of the most common cancers worldwide but is particularly prevalent in developed countries where it is associated with certain dietary factors and a sedentary lifestyle. One of the hallmarks of colorectal cancer cells is an altered metabolism that is not associated with tumor-causing mutations.

"Pyruvate is known to have an important role in <u>cancer metabolism</u>," explains lead author of the study Dr. Kenji Ohshima. "However, no one had previously studied how pyruvate produced by SRR is involved in cancer metabolism."

Therefore, the researchers set out to examine what role, if any, SRR plays in the development of colorectal cancer.

In a study published this month in *Nature Metabolism*, the researchers showed that SRR is much more abundant in colorectal cancer cells than in the surrounding tissues, and that <u>cancer cell lines</u> with higher levels of SRR divide at a much faster rate.

Says Dr. Ohshima, "This was a really exciting discovery because it showed that SRR is involved in colorectal cancer cell proliferation. When we then looked more closely at the pathway as a whole, we found that pyruvate levels were decreased in a slow-growing SRR mutant colorectal cancer cell line, suggesting that the pyruvate produced by SRR



enhances the proliferation of colorectal <u>cancer cells</u>. This result confirmed that the metabolic pathway is integral to the progression of colorectal cancer."

Given the obvious importance of SRR, the researchers then decided to test whether disruption of the <u>metabolic pathway</u> could prevent the progression of colorectal cancer. Astonishingly, not only did inhibition of SRR halt the growth of colorectal tumors in mice, it also improved the efficacy of currently available drugs used to treat colorectal cancer, causing a significant reduction in tumor size.

"There is still work to be done to confirm that our results translate into human cancer systems," explains senior author of the study Dr. Eiichi Morii. "However, based on these preliminary results, we expect that future strategies targeting SRR will provide effective new therapies for the treatment of colorectal <u>cancer</u>."

The article, "Serine racemase enhances growth of <u>colorectal cancer</u> by producing pyruvate from serine," was published in *Nature Metabolism*,

More information: Kenji Ohshima et al. Serine racemase enhances growth of colorectal cancer by producing pyruvate from serine, *Nature Metabolism* (2020). DOI: 10.1038/s42255-019-0156-2

Provided by Osaka University

Citation: Possible key to beating colorectal cancer (2020, January 29) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2020-01-key-colorectal-cancer.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.