

Toward a portable concussion detector that relies on an infrared laser

January 31 2020



Credit: CC0 Public Domain

With no way to reliably tell whether an athlete has a concussion, many may be playing with an undiagnosed injury. Likewise, 2 million people die every year because we don't have an early warning when brain cells



are dying—and another 4 million experience cognitive disabilities.

Now, a team of doctors and engineers has developed a noninvasive way to measure whether <u>brain</u> cells are in distress using an infrared laser. Their device uses optical fibers to deliver pulses of infrared light to a person's forehead. This light can penetrate through the skin and skull to the brain without doing harm. It interacts with an important molecule for metabolism called cytochrome C oxidase, or CCO.

While existing technology can give <u>real-time information</u> on whether the brain is getting oxygen, this new device gives information about whether the brain cells are able to use that oxygen. This information could enable concussions to be diagnosed on the sidelines of an athletic event. Alternatively, it could provide rapid feedback as doctors adjust treatment in the ER, operating room or the intensive care unit.

Mohammed Islam, professor of electrical engineering and computer science, will present the study on Super Bowl Sunday at the Photonics West conference in San Francisco. He and his collaborators at Michigan Medicine, the School of Kinesiology and the College of Literature, Science, and the Arts told us about how the new technique works.

What kinds of patients could this help?

Steven Broglio, director of the Michigan Concussion Center and professor athletic training: Concussions in athletics have been in the spotlight for the previous decade, but soldiers are also at high risk. Concussion can happen to anyone.

At present, there is no one test that can diagnose a concussion. There's no scan, no <u>blood test</u>, no computer test. We currently rely on a clinical exam and interpretation by the medical provider, so the diagnosis is subjective. We believe this technology is the first step in changing that



by measuring brain metabolism, potentially resulting in faster and more certain diagnoses.

The quicker you identify a concussion, the faster you can begin the recovery process of resting and gradually returning to normal activity. Research in the last 10 years has shown that resting sooner can shorten recovery time, helping people get back to school, work and military duty.

Rachel Russo, a trauma and emergency surgeon and clinical lecturer: This device could guide the treatment of more serious brain injuries and other conditions that put the brain at risk. In its broadest application, it could help patients whose blood pressure falls so much that there isn't enough blood flow to keep cells functioning normally. This can happen when a person is very ill or because they have lost too much blood. We need to keep them in the 'Goldilocks zone'—their <u>blood pressure</u> has to be high enough to supply oxygen to the cells and low enough to discourage bleeding in injured tissues.

In the ER and the intensive care unit, or ICU, patients are sometimes kept in medical comas while they heal from traumatic brain injuries. This would be the first device that could measure how well their <u>brain</u> <u>cells</u> are doing right in the hospital room.

What does measuring cytochrome C oxidase tell you?

Ioulia Kovelman, associate professor of psychology: It can tell us if the tissue is healthy and is metabolizing or "eating" properly—such as a brain of a healthy individual versus the brain of a person who suffered a concussion or has Alzheimer's disease, amyotrophic lateral sclerosis (ALS) or other types of brain dysfunction or trauma.

Russo: CCO decreases when cells are in distress. One of the challenges



is that we don't know if there's a universal threshold that signals a problem, the way we have normal ranges for different markers when we run a blood count. This is one of the areas we want to explore in the future.

But even without a threshold for action, we believe we could observe how a patient's cell metabolism changes over time and correlate that with other vital signs. This could give us an early indication of how they are recovering and whether more aggressive treatment is needed.

How does the device measure CCO?

Islam: There is a part of CCO that is excited by light with wavelengths between 750 nanometers and 900 nanometers. If you pluck a string on a violin, you hear its resonant frequency. CCO has, if you like, a string that we pluck with our laser, and we see the resonant frequency in the light we record.

Why hasn't it been done before?

Islam: Previous experiments used infrared lamps, but the signal from the CCO is not very distinct. Hemoglobin, which tells us about the blood oxygen levels, also responds to the range of wavelengths we use to probe CCO, and it is 10 to 20 times more plentiful in human tissue. We needed a stronger signal from the CCO to distinguish it from the hemoglobin and other molecules in the body, and we got that stronger response by using a laser that is nearly 10 times brighter than the lamps.

That increased brightness, along with measurement techniques to suppress noise during the measurements, was critical to enabling the conclusive measurements of CCO during cognitive attention tests with a portable, noninvasive device. Such attention tests are often used with



concussion patients.

What did your experiments show?

Kovelman: In our study, we asked participants to complete an attention task developed by Daniel Weissman, a professor of psychology, cognition and cognitive neuroscience. This task can be used to study healthy attention, such as attention development in children or attention dysfunction in individuals who might have a brain injury or concussion.

As individuals were completing this attention task, we saw an increase in oxygenated blood flow to their frontal lobe—the part of the brain that sits behind our forehead and is essential to being able to attend. Along with the increase in oxygenated blood flow, we also saw a change in the brain's CCO redox state—in other words, as oxygen supply increased, so did the consumption of the oxygen by the neurons of the frontal lobe that were busy attending to the task. This means that our new method can effectively measure brain metabolism, an index of brain function.

All cells use oxygen and energy, not just in the brain. Can this device look for trouble in other organs?

Russo: We think it can. We would have to adjust the depth of the measurement, but we may be able to observe other organs at risk of failure—for example, the heart, lungs or kidneys.

Provided by University of Michigan

Citation: Toward a portable concussion detector that relies on an infrared laser (2020, January 31) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2020-01-portable-concussion-detector-infrared-laser.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.