

Specialist technique used to study drugs completed for first time during Ebola outbreak

January 15 2020



Credit: CC0 Public Domain

A specialist technique used to study drugs has been completed for the first time during an outbreak of Ebola virus disease.

The study published today in *eBiomedicine* was a collaboration of researchers from Sierra Leone and the University's of Glasgow, Oxford, Cambridge and the Liverpool School of Tropical Medicine. It used pharmacokinetics—the measurement of the change in [drug concentration](#) in a person over time—to study an [experimental drug](#) called TKM-130803 for the treatment of Ebola [virus](#) disease during the 2015 outbreak in Sierra Leone.

Although treatment with TKM-130803 did not appear to improve survival, possibly because the patients presented with advanced disease, the research team were able to use pharmacokinetic analyses to gain important new insights.

In the study, the researchers measured the concentration of the drug over time in patients with Ebola. The scientists looked at the relationship between drug and Ebola virus concentrations and used the data to conduct computer simulations to answer important questions about what would happen if the dose and timing of drug administration were altered.

Significant findings were that the amount of virus (the 'viral load') was not significantly different in patients who lived or died, and that the concentration of drug was higher in those who died than in those who survived.

Dr. Janet Scott, Clinical Lecturer in Infectious Diseases at the University of Glasgow-MRC Centre for Virus Research, said: "The field team successfully gathered data and samples under extremely difficult conditions, right in the middle of the largest ever outbreak of Ebola Virus Disease in 2015. Rather than focus only on whether we could show that the drug worked or not, we were also able to study how the drug concentrations changed with time in patients. With this we were able to make a mathematical model, and predict what would probably have happened if we had used another dose.

"We had worried, that the dose chosen for this study might have been too low, but this analysis was able to show us that our caution had been justified, and that pushing the dose even higher would probably not have been safe. This reminds us that caution is important, especially when treating these extremely unwell patients."

Dr. Raman Sharma, senior post-doctoral research associate, Liverpool School of Tropical Medicine, said: "We have been able to show that even with data from very few patients, this kind of research is worth that extra effort, and can yield useful and important results, pushing our understanding of the drug and the dose beyond what can be learnt in the laboratory."

The laboratory analysis inside the Ebola Treatment Unit was led by Dr. Luc Meredith Senior Post Doctoral Scientist, University of Cambridge. He said: "It was difficult to remain accurate and keep the team safe, particularly in the excessive heat of northern Sierra Leone—but this needs to be done if we are to learn everything we can about Ebola and whatever new virus emerges next."

Prof. Peter Horby, Professor of Emerging Infectious Diseases and Global Health, University of Oxford said: "An important lesson is that pharmacokinetic analyses are possible even in very challenging settings and such analyses can provide important insights. This may be especially valuable in small or very quick outbreaks where there may be too few patients enrolled in trials to provide definitive proof of clinical impact. In such circumstance we can still measure the relationship between drug concentrations and virus concentrations, or other markers of disease progression, to give us an idea if the dose is adequate and drug is having an effect."

The study, "Pharmacokinetics of TKM-130803 in Ebola virus disease in Sierra Leonean: patients showed plasma concentrations which exceeded

target levels, with accumulation of [drug](#) in patients with most severe [disease](#)," is published in *eBiomedicine*.

More information: Janet T. Scott et al. Pharmacokinetics of TKM-130803 in Sierra Leonean patients with Ebola virus disease: plasma concentrations exceed target levels, with drug accumulation in the most severe patients, *EBioMedicine* (2020). [DOI: 10.1016/j.ebiom.2019.102601](#)

Provided by University of Glasgow

Citation: Specialist technique used to study drugs completed for first time during Ebola outbreak (2020, January 15) retrieved 4 May 2024 from <https://medicalxpress.com/news/2020-01-specialist-technique-drugs-ebola-outbreak.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--