

Technique is almost 86 percent effective in preventing maternal death from hemorrhaging

January 6 2020



Credit: CC0 Public Domain

According to the World Health Organization's latest statistics, from 2017, more than 800 women around the world die every day from preventable causes related to pregnancy and childbirth—the vast

majority of them in resource-poor areas. The leading reason is postpartum hemorrhage (PPH), which accounts for 27 percent of maternal mortality and occurs to some degree in five to seven percent of all deliveries.

Advanced interventions to stop the bleeding before the problem becomes unmanageable include emergency hysterectomy, a particular suturing technique, and ligation or embolization of the uterine artery—but because these require extensive training and/or high-tech equipment, not all hospitals around the world have access to them. A simple, inexpensive alternative, the uterine balloon tamponade (UBT), has been available since the early 1980s. In it, a balloon is inserted through the cervix by catheter and inflated with water, filling the [uterine cavity](#) and arresting the hemorrhage.

Two studies—published in 2013 and 2019—brought into question the effectiveness of UBT, but a paper published today in the *American Journal of Obstetrics and Gynecology* "really puts a nail in that coffin," according to principal investigator Thomas Burke, MD, FACEP, FRSM, director of the Global Health Innovation Lab in the Emergency Department at Massachusetts General Hospital and an associate professor at Harvard Medical School and the Harvard T.H. Chan School of Public Health.

"The reason we undertook this study is that those two trials caused a great deal of confusion and controversy regarding UBT," Burke said. "We decided to take an extremely rigorous approach, bringing together all of the world's literature on this technique and inviting independent investigators to join. It was a massive project." Burke and his colleagues conducted a [systematic review](#) and meta-analysis of 91 randomized control trials, nonrandomized studies, and case series, and found an overall success rate of 85.9 percent for UBT.

"So what this really teaches us is that the challenges experienced in these two trials were about the program implementation, not the device itself," he said. "There's no question the device works fine. But saving a life is much more complex than just handing someone a device that's new to their practice. We need to study how to integrate the device into a health system so that good uptake, appropriate use, and best practice result in quality care."

Burke pointed out that it is "exquisitely clear" that in many areas of the world, maternal mortality drops significantly when UBT is used. "They have saved many lives," he said. "But there are still places where one in six women lose their lives from pregnancy-related causes and in 2020 that should never be. It's an embarrassment to humankind."

More information: Sebastian Suarez et al, Uterine Balloon Tamponade for the Treatment of Postpartum Hemorrhage: a Systematic Review and Meta-Analysis, *American Journal of Obstetrics and Gynecology* (2020). [DOI: 10.1016/j.ajog.2019.11.1287](https://doi.org/10.1016/j.ajog.2019.11.1287)

Provided by Massachusetts General Hospital

Citation: Technique is almost 86 percent effective in preventing maternal death from hemorrhaging (2020, January 6) retrieved 3 May 2024 from <https://medicalxpress.com/news/2020-01-technique-percent-effective-maternal-death.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--