


# Shining new light on the actions of key neurotransmitter in the brain

February 25 2020

---



GABA neurotransmitter. Credit: Florey Institute of Neuroscience and Mental Health

An international research collaboration has uncovered evidence that a common neurotransmitter can selectively regulate the excitability of neurons.

The researchers were investigating [gamma-aminobutyric acid](#), known as GABA, the main inhibitory neurotransmitter in the human [brain](#) when the findings were made.

"We were using computer models from the Blue Brain Project which predicted that GABA could be having two different functions—increasing the excitability of one type of interneuron, and decreasing the excitability of another type of interneuron," said Dr. Alexander Bryson from the Florey Institute of Neuroscience and Mental Health.

"This was surprising to us because GABA is primarily thought to inhibit or reduce the excitability of neurons," said Dr. Bryson.

The researchers were subsequently able to observe the phenomenon in the laboratory. These results challenge the prevailing scientific view and suggest that sub-types of interneurons defined by their electrophysiology characteristics can undergo selective modulation by GABA.

Professor Sean Hill, co-Director at the EPFL Blue Brain Project says the results were only possible through collaboration bringing together the modelling expertise and computational infrastructure of Blue Brain, with the receptor expertise and data from the Florey Institute.

"The result was wholly counterintuitive and exciting, with significant implications for understanding brain circuit alterations in [mental health disorders](#)," said Professor Hill.

Professor Steven Petrou, Director of the Florey Institute agrees that the finding has exciting implications.

"Understanding the [human brain](#) is one of the great challenges remaining for scientists. GABA is the predominant chemical messenger in the brain and we already knew its role changes over the course of neurodevelopment. These findings show additional complexity of this common neurotransmitter," said Professor Petrou.

The research has been published in *Proceedings of the National Academy of Sciences*.

**More information:** Alexander Bryson et al. GABA-mediated tonic inhibition differentially modulates gain in functional subtypes of cortical interneurons, *Proceedings of the National Academy of Sciences* (2020). [DOI: 10.1073/pnas.1906369117](https://doi.org/10.1073/pnas.1906369117)

Provided by Florey Institute of Neuroscience and Mental Health

Citation: Shining new light on the actions of key neurotransmitter in the brain (2020, February 25) retrieved 4 May 2024 from <https://medicalxpress.com/news/2020-02-actions-key-neurotransmitter-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--