

Autism eye scan could lead to early detection

February 21 2020



Dr. Paul Constable using eye scan device to potentially detect autism Credit: Flinders University

A new eye scan could help identify autism in children years earlier than currently possible.

The non-invasive eye scan utilises a hand-held device to find a pattern of

subtle electrical signals in the retina that are different in [children](#) on the [autism](#) spectrum, which are directly linked to differences in their brain development.

The scan was tested on about 180 people with and without autism between the ages of 5 and 21 in collaboration with Yale University in the US, University College London and Great Ormond Street Hospital in the UK, as part of a study published in the *Journal of Autism and Developmental Disorders*.

These potential biomarkers for [autism spectrum](#) disorder (ASD) could allow for early detection of other disorders, such as attention deficit hyperactivity disorder (ADHD).

Dr. Paul Constable, a senior lecturer at the College of Nursing and Health Sciences at Flinders University, has been searching for an autism "biomarker" since 2006, in an effort to improve early detection and intervention methods after his own child was diagnosed.

"The retina is an extension of the brain, made of neural tissue and connected to the brain by the optic nerve, so it was an ideal place to look," says Dr. Constable.

"The test is a quick, non-intrusive eye-scan using a hand-held device and we anticipate it will be equally effective on [younger children](#). Very early diagnosis means not only can children receive important interventions, but families are empowered to get the necessary supports in place, come to terms with the diagnosis, and make informed decisions."

Dr. Constable's research team is also investigating the scan to detect autism in younger children and other conditions including attention deficit hyperactivity disorder, and other neurodevelopmental [disorders](#).

"Now we have found a likely candidate biomarker for autism, the next stage is to look at young children, even infants, as the earlier we can get to intervention stages the better," Dr. Constable says.

He says his team often encounters parents who have two or three young children with autism, as the chance of having a second autistic child is much higher for parents with one child on the spectrum. Autism in Australia is usually diagnosed after the age of four.

Early detection in firstborn children could give parents the opportunity to decide if they want to have more kids, with previous studies indicating an increased likelihood siblings of autistic children are more likely to develop the disorder.

"Detection inevitably changes family dynamics and goals, and creates consideration about the time required to help the child," Dr. Constable says.

"Very [early diagnosis](#) means not only can children receive important interventions, but families are empowered to get the necessary supports in place, come to terms with the diagnosis, and make informed decisions."

More information: Paul A. Constable et al, Light-Adapted Electroretinogram Differences in Autism Spectrum Disorder, *Journal of Autism and Developmental Disorders* (2020). [DOI: 10.1007/s10803-020-04396-5](https://doi.org/10.1007/s10803-020-04396-5)

Provided by Flinders University

Citation: Autism eye scan could lead to early detection (2020, February 21) retrieved 13 March

2024 from <https://medicalxpress.com/news/2020-02-autism-eye-scan-early.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.