

# Got a cervix? Get it tested or test it yourself

February 4 2020

---



Gardasil vaccine and box. Image: Wikipedia

Women who have sex with women (WSW), transsexual men and non-binary people with a cervix are at similar risk of human papillomavirus (HPV) infection and subsequent risk of developing cervical cancers as heterosexual women, but are often never screened or underscreened, according to the authors of a Lessons from Practice article published online by the *Medical Journal of Australia*.

Dr. Monica McGauran and Dr. Adam Pendlebury, from the Mercy Hospital for Women in Melbourne, reported the case of a 39-year-old woman who was referred for colposcopy with a positive human papillomavirus (HPV) type 16 infection result on a self-collected HPV

vaginal swab. She was asymptomatic and was overdue by many years for [cervical screening](#).

"Barriers to regular cervical screening were cited as having difficulty with speculum examination and having a female partner, with the subsequent perception that cervical screening was not required," McGauran and Pendlebury wrote.

"HPV can be transmitted intravaginally between female sexual partners via digital contact and fomites, such as sex toys. In addition, a proportion of WSW have had previous or have current male sexual partners."

The authors cited 2001 research which found that 10% of WSW had been told by health care practitioners that they did not need to be screened if they did not have sex with men.

"Other reasons for reduced uptake [of screening] included lower perceived level of risk for HPV infection, provision of health care that was not sensitive or inclusive to non-[heterosexual women](#), and reduced provision of opportunistic health care due to the reduced need for contraception," wrote McGauran and Pendlebury.

Self-collected samples are now available as part of the National Cervical Screening Program for [women](#) over the age of 30 years who have never been screened or who are overdue for [screening](#) by two or more years. It is available to be collected in a practitioner-supervised setting (ie, not at home) and is not available to women who are pregnant, symptomatic, or who have a past history of dysplasia.

"Self-collected vaginal swabs for HPV polymerase chain reaction are as sensitive as practitioner-collected samples and present an excellent option for never-screened and underscreened women, transsexual men and non-binary people with a cervix who are aged 30 years and older and

who decline practitioner collection," McGauran and Pendlebury concluded.

**More information:** Monica FG McGauran et al. HPV swab self-collection and cervical cancer in women who have sex with women, *Medical Journal of Australia* (2019). [DOI: 10.5694/mja2.50457](https://doi.org/10.5694/mja2.50457)

Provided by University of Melbourne

Citation: Got a cervix? Get it tested or test it yourself (2020, February 4) retrieved 28 April 2024 from <https://medicalxpress.com/news/2020-02-cervix.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.