

What is Charles Bonnet syndrome, the eye condition that causes hallucinations?

February 5 2020, by Jason Yosar



Credit: Unsplash/CC0 Public Domain

Visual hallucinations, or seeing things that aren't really there, can be frightening and distressing.

They may occur due to a large variety of physical and psychiatric conditions. But a lesser known cause is Charles Bonnet syndrome (pronounced *bo-nay*), named after the [Swiss scientist](#) who first described the condition in 1760.

Charles Bonnet syndrome (also called visual release hallucinations) refers to visual hallucinations in patients with severe vision loss due to eye, optic nerve or [brain disease](#).

We don't know the exact cause of Charles Bonnet syndrome. But the most commonly accepted theory is the loss of visual sensory signals to the brain (for example, when a person becomes blind) means the brain cannot put the brakes on [excessive and unwanted brain activity](#).

This leads the part of the brain responsible for the sensation of vision (the visual cortex) to fire signals inappropriately. The person in turn perceives they are seeing something in the absence of a true stimulus—a visual hallucination.

If these symptoms are affecting you, a friend or family member who has become blind in one or both eyes, it's important to understand it's not a sign of "going mad."

What are Charles Bonnet hallucinations like?

The hallucinations may be "simple" (such as lines, shapes, or flashes of light) or "complex" (such as formed images of animals, like butterflies). Simple hallucinations are much more common.

They may occur for seconds or minutes to hours or continuously, and the [frequency ranges](#) from isolated episodes to multiple times a day. It's normal for Charles Bonnet syndrome to [last for years](#); some people will experience symptoms for the rest of their lives.

The nature of Charles Bonnet hallucinations is highly variable. That is, people who are affected often don't see the same thing repeatedly, and one person with Charles Bonnet syndrome will see different things from the next person.

Charles Bonnet hallucinations often have little or no emotional meaning, allowing affected people to [recognize they are not real](#). This is distinct from hallucinations associated with mental illness.

Other features of visual hallucinations unique to Charles Bonnet syndrome include:

- hallucinations only appear in the areas where vision is lost (for example, a person who is blind in their left eye will perceive hallucinations only in that eye)
- hallucinations are more frequently seen with the [eyes open than closed](#), and may disappear when the person closes their eyes or looks away
- hallucinations are more common in settings of [sensory deprivation](#) (for example, at night time or in dim lighting, or during periods of inactivity).

Who is affected?

Most people with Charles Bonnet syndrome are [older adults](#) (usually over 70). This is probably because vision loss is most common in this age group. But any person of any age with acquired vision loss can develop Charles Bonnet syndrome.

The causes of blindness that lead to Charles Bonnet syndrome are usually [macular degeneration](#), glaucoma, [diabetes](#), [stroke](#) and injury—but any disease that leads to blindness may cause Charles Bonnet syndrome.

The syndrome does not occur in congenital blindness (people born blind from birth).

We currently have no conclusive data on how many Australians have Charles Bonnet syndrome, although [one study](#) estimated more than 17% of people aged over 60 with impaired vision had it. In [another study](#), as many as 57% of participants with vision loss reported perceived visual hallucinations.

Importantly, it may be more common than estimated because of lack of reporting. That is, people who are affected may not report their hallucinations due to fear of psychiatric disease or of being perceived to be "going mad."

Further, people who do report their symptoms may be [misdiagnosed](#) with psychosis or dementia.

Treatment options are limited

Seeing a [general practitioner](#) (often in conjunction with a neurologist and/or geriatrician) is an important first step to exclude other causes of hallucinations. These could include dementia, physical neurological conditions (for example, a brain tumor), epilepsy and delirium due to infections or medications. Your doctor may order blood tests and/or brain imaging to rule these out.

Treatment for Charles Bonnet [syndrome](#) is very limited, but [many patients report](#) reassurance is all they need, especially for infrequent hallucinations or those that don't adversely affect quality of life.

Strategies to minimize the frequency and duration of [hallucinations](#) include frequent blinking or rapid eye movement, going to a lighter place or switching a light on, and increasing social interaction, which helps to

counter inactivity.

For patients with debilitating symptoms, doctors may trial medications such as [antidepressants](#), [antipsychotics](#) and [antiepileptic drugs](#), though their efficacy is variable and may be outweighed by side effects.

Hallucinations may disappear if the cause of [vision](#) loss can be corrected (for example, if severe cataracts were causing blindness and the patient has a cataracts operation).

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: What is Charles Bonnet syndrome, the eye condition that causes hallucinations? (2020, February 5) retrieved 25 April 2024 from <https://medicalxpress.com/news/2020-02-charles-bonnet-syndrome-eye-condition.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--