

Childhood physical abuse linked to heavy cigarette use among teens who smoke

February 24 2020



Credit: CC0 Public Domain

Researchers have known that kids who are at high risk of being mistreated at home—who live in poverty or have parents who use drugs or have mental health problems—are more likely to start smoking. Because abused and neglected children are often unsupervised, these

teens have easy access to cigarettes and other substances that they use to deal with anxiety and other trauma-related symptoms.

A new study shows that physical [abuse](#) of children in high-risk homes, especially when they're toddlers or teens, dramatically increases the odds that their adolescent experimentation with cigarettes will lead to a heavy smoking habit.

Neglect of very young children in the same high-risk population was linked to a gradual increase in [cigarette use](#) among teens who smoked.

The study examined data on children who were at high risk for abuse and neglect—either because they had been referred to a child protective service or lived in conditions associated with the likelihood of maltreatment, or both.

"I wanted to look at different types of maltreatment and whether they have an impact on cigarette smoking," said lead author Susan Yoon, assistant professor of social work at The Ohio State University. "There's also the impact of timing—it's not just maltreatment, yes or no. When did it happen, and does that matter with regard to outcomes?"

The study is published online in the journal *Substance Use and Misuse*.

Yoon noted that substance abuse research doesn't often take cigarette smoking into consideration despite its significant damaging effects on health.

"Adolescent cigarette smoking is a really serious social problem and public health concern. Brain development is not complete until late adolescence or during young adulthood, and cigarette smoking is associated with damage in [brain development](#)," Yoon said. "We also know that those who start smoking cigarettes during adolescence are

more likely to continue smoking into adulthood."

Yoon and colleagues used data from the Longitudinal Studies of Child Abuse and Neglect, a consortium of studies across different regions of the United States targeting children who were identified as maltreated or at risk for maltreatment. This study used data on 903 adolescents who were assessed at age 12, 16 and 18.

A breakdown of different types of abuse and neglect experienced by the sample population during three different time periods ([early childhood](#), school age and adolescence) confirmed how vulnerable these kids were. Almost a quarter had been neglected during early childhood and their school-age years, and 19 percent reported school-age physical abuse. Almost half lived in poverty and 58 percent lived in households in which family members smoked cigarettes.

Yoon used their responses about smoking between the ages of 12 and 18 to identify three patterns of cigarette use: stable low/no use (61 percent of respondents), gradually increasing use (30 percent) and sharply increasing cigarette use (9 percent).

"It was almost shocking how the pattern of cigarette use over time went up so drastically in the sharply increasing use class," Yoon said. "They were pretty similar to the others at age 12—almost 80 percent didn't smoke. At age 16, we saw that almost 60 percent had used cigarettes more than 20 days in the past year and by 18, every single kid in this group reported heavy use of cigarettes."

A [statistical analysis](#) showed that adolescents who experienced early childhood physical abuse were 2.3 times more likely to be in the sharply increasing cigarette use group compared with the stable no/low group. Physical abuse during adolescence had an even stronger effect—this type of mistreatment at that point in life was linked to 3.7 times higher

odds for sharply increased cigarette use.

Adolescents who had been neglected during early childhood were 1.89 times more likely to be in the gradually increasing cigarette use group than in the stable no/low use group. Neglect can include a lack of appropriate supervision and failure to meet kids' needs—ranging from food and clothing to medical needs and education.

About 40 percent of these smokers had reported using cigarettes at age 16, and by age 18, more than 80 percent were smokers, and about 40 percent had smoked on more than 20 days in the previous year.

The analysis also measured the potential effects of sexual abuse and emotional abuse on cigarette use, but neither was a predictor of an increase in smoking.

"The different types of maltreatment predicted different patterns of cigarette use, but the timing mattered as well. It was early childhood and adolescence that appear to be really sensitive periods in which maltreatment influences these outcomes," Yoon said.

The results suggest efforts to prevent [smoking](#) by teens at risk of maltreatment—and especially children experiencing physical abuse and neglect—should start before age 12, she said.

"If we intervene too late, it's likely they've already started using cigarettes and are in the process of using more cigarettes," Yoon said.

"We know from our studies that these kids experiencing physical abuse and neglect are at much higher risk of showing this poor pattern of cigarette use over time. We should be targeting this high-risk group for cigarette use prevention and intervention."

More information: Susan Yoon et al, The Type and Timing of Child

Maltreatment as Predictors of Adolescent Cigarette Smoking Trajectories, *Substance Use & Misuse* (2020). DOI: [10.1080/10826084.2020.1713819](https://doi.org/10.1080/10826084.2020.1713819)

Provided by The Ohio State University

Citation: Childhood physical abuse linked to heavy cigarette use among teens who smoke (2020, February 24) retrieved 24 April 2024 from <https://medicalxpress.com/news/2020-02-childhood-physical-abuse-linked-heavy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.