

# New study adds to evidence of diabetes drug link to heart problems

February 5 2020

---

A new study published by *The BMJ* today adds to evidence that rosiglitazone—a drug used to treat type 2 diabetes—is associated with an increased risk of heart problems, especially heart failure.

This study is the most comprehensive evaluation of the cardiovascular risk of [rosiglitazone](#) ever done.

Rosiglitazone belongs to a class of drugs called thiazolidinediones. It helps control blood sugar levels in patients with type 2 diabetes, but it can also increase the risk of serious heart problems. This has led to suspension of the drug in Europe and previous restrictions on its use in the United States.

However, since 2007, studies have reported conflicting findings about whether rosiglitazone increases the risk of heart attacks. But these studies didn't have access to the [raw data](#), also known as individual patient level data (IPD), from [clinical trials](#) and mostly relied on summary level data (e.g. results reported in publications and clinical trial registries), which are not as reliable when estimating the true safety profile of drugs.

Recent efforts by GlaxoSmithKline (GSK) - the maker of rosiglitazone—to make IPD available to external investigators, prompted a team of US researchers to re-analyse the data and clarify some of the uncertainties about rosiglitazone's cardiovascular risk.

They analysed the results of more than 130 [trials](#) involving over 48,000 [adult patients](#) that compared rosiglitazone with any control for at least 24 weeks. IPD were available for 33 trials, which included 21 156 patients; the remaining trials only had summary level data available.

When the researchers analysed the IPD from trials made available by GSK, they found rosiglitazone was associated with a 33% increased risk of a composite cardiovascular event (heart attack, [heart failure](#), cardiovascular and non-cardiovascular related death) compared with controls. This was estimated from the 274 events among 11,837 rosiglitazone patients and 219 events among 9,319 control patients.

When examining cardiovascular events independently, the analyses of the 33 GSK trials with IPD resulted in higher estimates of the risk of [heart](#) attacks than the analyses of trials with IPD and summary level data.

These findings highlight the potential for different results derived from different data sources, and demonstrate the need for greater clinical trial transparency and data sharing to accurately assess the safety of drugs, say the researchers.

"Our study suggests that when evaluating drug safety and performing meta-analyses focused on safety, IPD might be necessary to accurately classify all adverse events," they write. "By including these data in research, patients, clinicians, and researchers would be able to make more informed decisions about the safety of interventions."

They add: "Our study highlights the need for independent evidence assessment to promote transparency and ensure confidence in approved therapeutics, and postmarket surveillance that tracks known and unknown risks and benefits."

**More information:** Updating insights into rosiglitazone and

cardiovascular risk through shared data: individual patient and summary level meta-analyses, *BMJ* (2020). DOI: [10.1136/bmj.17078](https://doi.org/10.1136/bmj.17078)

Provided by British Medical Journal

Citation: New study adds to evidence of diabetes drug link to heart problems (2020, February 5) retrieved 13 March 2024 from <https://medicalxpress.com/news/2020-02-evidence-diabetes-drug-link-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.