

# Further evidence adds weight to call for urgent action on salt targets

February 25 2020

---



Credit: CC0 Public Domain

Most people get a blood pressure lowering benefit from eating less salt and the less you eat, the more you benefit, but for older people and those who already have high blood pressure the effect is even greater,

according to new research published in the *British Medical Journal* today.

The study also suggests that if you reduce your [salt](#) intake for longer, you're likely to see a larger fall in your blood pressure.

George Institute for Global Health researchers, as part of an international collaboration, conducted the most up to date [systematic review](#) and meta-analysis of randomized controlled trials that examined the effect of reducing salt intake on blood pressure.

Lead author Polly Huang, Ph.D. candidate and data analyst at The George Institute said that while there was consensus among health and scientific organizations about reducing salt intake in the [general population](#), some have proposed that it has little value for those with normal blood pressure.

"Our study showed the benefits of lowering salt intake are seen across the board, but if you're older and already have high blood pressure—and are therefore at greater risk of associated conditions such as stroke and heart disease—there is an even greater effect on blood pressure from the same amount of salt reduction," she said.

High blood pressure is one of the biggest contributors to premature death from stroke or heart disease and too much salt in the diet is one of the major culprits. At about 9g per day, average salt intake in Australia is almost double WHO recommendations (5g per day). Worldwide, excess salt intake is estimated to cause about three million deaths each year.

Ms Huang said that the effects of salt reduction were more evident at higher blood pressure levels, older ages, and among non-white populations, but almost every population group examined achieved a reduction in blood pressure.

"More importantly, salt reduction in people with normal blood pressure could potentially prevent or delay the development of [high blood pressure](#) with age," she added.

The George Institute's Senior Public Health Nutritionist Clare Farrand said the findings added more weight to the argument for reducing population-wide salt intake as a key public health strategy to lower blood pressure and reduce the risk of strokes and [heart disease](#).

"The majority of the salt we eat is hidden in processed and packaged foods and most people aren't aware that the amount they are consuming is raising their [blood](#) pressure and shortening their lives," she said.

"Setting salt targets for the [food industry](#) works, but Australia has been lagging behind other countries on this. We're pleased to now see the Government's Healthy Food Partnership endorsement of a 'first wave' of salt targets, with a second expected to follow mid-2020."

"However, to meet the stated commitment to a 30 percent reduction in the average population [salt intake](#) by 2025, the targets need to be rapidly implemented and food companies closely monitored to ensure they're meeting these targets," added Ms Farrand.

"With more than six million Australian adults—that's more than a third of our adult population, or one in three people—having high [blood pressure](#), reducing Australian salt consumption would save thousands of lives each year as well as millions in healthcare costs."

**More information:** Liping Huang et al. Effect of dose and duration of reduction in dietary sodium on blood pressure levels: systematic review and meta-analysis of randomised trials, *BMJ* (2020). [DOI: 10.1136/bmj.m315](#)

Provided by George Institute for Global Health

Citation: Further evidence adds weight to call for urgent action on salt targets (2020, February 25) retrieved 2 May 2024 from <https://medicalxpress.com/news/2020-02-evidence-weight-urgent-action-salt.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.