

# Ibuprofen might make your periods lighter, but it's not a long-term solution

February 10 2020, by Mia Schaumberg

---



Credit: AI-generated image ([disclaimer](#))

A tweet saying ibuprofen reduces menstrual flow by 50% went viral last month.

The original tweet and ensuing responses fueled a debate about society's supposed unwillingness to talk about periods.

At the same time, many [women](#) responded with questions about the evidence behind this notion, how it works, and the risks.

There's [some evidence](#) non-steroidal anti-inflammatory drugs (NSAIDs), like [ibuprofen](#), can reduce [menstrual flow](#). But they shouldn't be seen as a long-term solution. Women who experience heavy or painful periods regularly should speak to a doctor.

## How does this work?

Menstrual symptoms, including heavy periods, affect quality of life for many women. Heavy menstrual bleeding is one of the most common reasons women visit a gynecologist, accounting for [up to 30% of visits](#).

One study found menstrual symptoms including pain, heavy bleeding and low mood may be linked to nearly [nine days](#) of lost productivity per woman every year.

Women who have heavy bleeding, and who experience painful periods, have elevated levels of hormones called [prostaglandins](#). Prostaglandins act to dilate [blood vessels](#), slow the clotting process and also help the body shed the uterine lining by inducing muscle contractions. So having higher levels of these hormones can lead to heavier bleeding and more severe cramping.

Ibuprofen has been shown to [reduce prostaglandin levels](#) in the lining of the uterus, which may be one way it reduces menstrual flow, though the exact mechanism remains uncertain.

This process may also be part of the reason ibuprofen can be an effective [first-line treatment](#) option for painful periods.

## What the evidence says

A [recent review](#) of the research on this topic found on the whole, NSAIDs were more effective than a placebo at reducing [blood loss](#) in women with heavy menstrual bleeding.

But only [one study](#) in this review compared ibuprofen specifically to a placebo. This study, published in 1986, included 24 women. Half were given ibuprofen, and half a placebo. There was a modest reduction of 36mL (25%) in menstrual blood flow with ibuprofen treatment. This study is obviously very small, so doesn't provide evidence we'd consider strong.

So, the evidence doesn't support the 50% reduction in menstrual flow the tweeter claimed.

The review found another drug, tranexamic acid, was actually more effective than NSAIDs at reducing menstrual flow, [with a 54% reduction in menstrual blood loss](#). However, it isn't available over the counter, making it less accessible.

It's also important to note this review looked at women with heavy periods. There's no strong evidence to suggest ibuprofen, or other NSAIDs, can significantly reduce menstrual flow in women with regular, healthy menstruation.

According to the review, ibuprofen also does not appear to be effective at reducing menstrual flow in women where a pre-existing [medical condition](#) is the reason for heavy bleeding. These conditions [might include](#) uterine fibroids (non-cancerous growths in the uterus walls), clotting disorders or hormonal imbalances associated with conditions such as polycystic ovary syndrome.

## Are there any risks to consider?

Ibuprofen may provide some relief in the short term, and it may offer a level of convenience in being available over the counter, but it's generally not recommended as a long-term treatment for heavy periods.

[Long term use of NSAIDs](#) such as ibuprofen has been linked with kidney disease, blood pressure issues and stomach ulcers. Other common side effects include indigestion, headaches and drowsiness, especially when it's taken in higher doses.

Using ibuprofen may not be suitable for people with existing conditions such as liver or kidney disease or stomach ulcers.

Ibuprofen or other NSAIDs should only be used as a first-line treatment, before discussing longer-term solutions with a medical practitioner.

If heavy and/or painful periods are a consistent issue, there might be an underlying cause. In this case, the evidence suggests ibuprofen does not significantly reduce menstrual flow. So if this is something you struggle with every month, talk to your doctor to find a safe, long-term solution.

There are many evidence-based options available for managing heavy menstrual bleeding in the longer term, such as the oral contraceptive pill or the [hormonal IUD](#). Your doctor can assess your individual circumstances and potential risk factors to see what will be right for you.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Ibuprofen might make your periods lighter, but it's not a long-term solution (2020, February 10) retrieved 24 April 2024 from <https://medicalxpress.com/news/2020-02-ibuprofen-periods-lighter-long-term-solution.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.