

# **IVF-conceived children have somewhat higher mortality risk in their first weeks of life**

February 18 2020

---



Credit: CC0 Public Domain

Children conceived with assisted reproductive techniques including in vitro fertilization (IVF) have a somewhat higher mortality risk during

their first weeks of life than children conceived naturally, according to a study by researchers at Karolinska Institutet in Sweden published in the journal *Fertility and Sterility*. The researchers link the increased risk to a higher degree of premature births in IVF children and emphasize that the risk of infant mortality is still very small for both groups. Beyond one year of age, the risk of mortality was similar for all children regardless of conception method.

"It is important to note that even if we on a group level can see a somewhat increased risk of infant mortality after IVF, the absolute risk for each individual is still very small," says Kenny Rodriguez-Wallberg, associate professor at the Department of Oncology and Pathology at Karolinska Institutet and the corresponding author. "It is also reassuring to know that there is no increased risk of mortality in this group of [children](#) beyond the first year of life."

More and more women seek help to become pregnant and the achieving of pregnancies made possible with assisted reproductive techniques have increased in recent years. In most cases, these pregnancies have a happy outcome with a healthy baby. Prior studies show, however, that IVF-pregnancies come with an [increased risk](#) of low birth weight, prematurity and birth defects. These risks have partly been linked to the increased probability of twin-births after IVF-treatment.

In the current study, the researchers selected only singleton children and compared mortality in children conceived through different types of assisted reproductive techniques with children who were conceived naturally. They analyzed data on 2.8 million children born in Sweden over a period of 30 years. Some 43,500 of these were the result of assisted reproduction.

In total, 7,236 children died before one year of age, of whom only 114 were conceived with assisted reproductive techniques. After adjusting

for confounding factors such as the mother's age and earlier infertility, the researchers found that the children conceived through IVF had a 45 percent higher risk of death before one year of age than children conceived naturally. The level of risk varied depending on which type of assisted reproductive technique was used, and how many days had passed since birth. The risk gradually declined after the first weeks of life.

During the first week of life, the children conceived after transfer of a frozen embryo had a more than two-fold higher risk of death than the children conceived naturally. This was, however, based on only a small sample of children conceived with frozen embryos. After one week, the risk dropped to about the same level as the naturally conceived children. Infants conceived from transfer of a fresh embryo or with the help of an intracytoplasmic sperm injection (ICSI)—where a single sperm is injected into the egg—did not have a higher risk of death than naturally conceived children, irrespective of how many days had passed since birth.

"Our results indicate that the kind of assisted reproductive technique used may make a difference, and therefore it is important to further investigate what causes or underlying mechanisms are behind the risks," says Anastasia Nyman Iliadou, associate professor at the Department of Medical Epidemiology and Biostatistics at Karolinska Institutet and senior lead author. "They also show the need for extra attention and care of children conceived with IVF, especially during the first week of life."

According to the researchers, one explanation may be that more IVF-children are born prematurely than those conceived naturally, which in itself could have negative consequences. It is also possible that the underlying infertility leads to a higher risk of complications. The leading causes of infant [mortality](#) among children conceived with assisted reproductive techniques included [respiratory distress](#), incomplete lung

development, infections and neonatal hemorrhage, which are conditions often linked to prematurity.

**More information:** "Mortality from infancy to adolescence in singleton children conceived from assisted reproductive techniques (ART) vs naturally conceived singletons in Sweden," Kenny A. Rodriguez-Wallberg, Frida E. Lundberg, Sara Ekberg, Anna L. V. Johansson, Jonas F. Ludvigsson, Catharina Almqvist, Sven Cnattingius and Anastasia N. Iliadou, *Fertility and Sterility*, online Feb. 18, 2020.

Provided by Karolinska Institutet

Citation: IVF-conceived children have somewhat higher mortality risk in their first weeks of life (2020, February 18) retrieved 3 May 2024 from <https://medicalxpress.com/news/2020-02-ivf-conceived-children-higher-mortality-weeks.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--