


Language disorders as indicators of the diagnosis and progression of Huntington's disease

February 20 2020



HTT gene neuron. Credit: <https://commons.wikimedia.org/wiki>

Huntington's disease is a hereditary neurodegenerative disorder caused by a gene on chromosome 4 that affects a very important area of the brain, the striatum. People are born with the defective gene but symptoms do not appear until the age of 30 or 40. This disease, in

addition to motor impairments, cognitive and affective problems, also involves changes in language. A study shows that the first symptoms of the disease are revealed through linguistic changes in spontaneous speech.

This is revealed in a study published in *Journal of Communication Disorders* by Antonia Tovar as first author, led by Wolfram Hinzen, ICREA research professor, both researchers with the Department of Translation and Language Sciences at UPF, involving researchers from the universities of Paris-Saclay (Orsay, France), the UAB, the UB, together with the Hospital Mare de Déu de la Mercè in Barcelona and FIDMAG Sisters Hospitallers Research Foundation (Barcelona).


"Linguistic impairments in Huntington's disease arise before the onset of motor impairment and even when neuropsychological tests prove normal. This suggests that [language](#) may be a biomarker for the progression of Huntington's," Tovar and Hinzen explain.

Specific language patterns of Huntington's disease

In addition to showing that linguistic impairments in Huntington's disease arise before the onset of motor impairment, the authors have sought to characterize the nature of these specific changes in Huntington's disease. Thus "language needs to be assessed as a multi-dimensional construct organized at multiple levels (phonology, morphology, syntax, semantics, discourse), which are re-fitted together into an integrated functional whole," the authors point out.

The few existing linguistic studies on spontaneous speech in Huntington's disease have reported language shortcomings. For example patterns of reduced syntactic complexity, with fewer words and syntactic structures formed in short, simple sentence constructions, more paraphasic and grammatical errors, etc. These language deficits may

form a distinctive signature profile of Huntington's as compared with Parkinson's disease or people with cerebral lesions.



Boxplots for differences between groups in the analysis of linguistic variables grouped into five linguistic domains. Credit: UPF

To study the language profiles of Huntington's disease, the authors designed an experiment with 20 HD gene-carriers, all native Spanish speakers; 10 in the early stages of the disease and another 10 with pre-manifest Huntington's disease, and their respective neurotypical controls not carrying the gene, and compared them regarding the structure and function of language in spontaneous speech. The researchers also studied the relationship between language and non-linguistic cognitive impairment.

"This disease is generally accepted to begin with the onset of motor symptoms, but we have found differences in the language profile of subjects without motor symptoms," says Tovar, first author of the article.

In addition to proving that in Huntington's disease linguistic changes precede other cognitive and motor impairments, it has been shown that language impairments are in the areas of basic grammar organization and cannot be detected with standard neurocognitive tests. This study is important because it has managed to capture subtle changes in the patient's linguistic profile that could not be detected using the tests generally conducted in this population.

A more detailed and differentiated profile of the linguistic phenotype of this disease

The results showed that while the patients mentioned have "pauses" in their speech, at the beginning of Huntington's disease they tend to be filled through prolongations and repetitions. Therefore, the study provides further support for the disintegration of language in Huntington's disease and contributes to a more detailed and

differentiated profile of the linguistic phenotype of the disease.

Since [language skills](#) mainly have an impact on [communication skills](#), the authors conclude that special attention must be paid to the early detection of Huntington's disease by means of clinical linguistic tests, and the sensitivity and specificity of language as a marker for the progression of the disease must continue to be systematically investigated.

More information: Antonia Tovar et al, Language disintegration in spontaneous speech in Huntington's disease: a more fine-grained analysis, *Journal of Communication Disorders* (2019). [DOI: 10.1016/j.jcomdis.2019.105970](#)

Provided by Universitat Pompeu Fabra - Barcelona

Citation: Language disorders as indicators of the diagnosis and progression of Huntington's disease (2020, February 20) retrieved 3 May 2024 from <https://medicalxpress.com/news/2020-02-language-disorders-indicators-diagnosis-huntington.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--