


MRI findings predict shoulder stiffness for rotator cuff tears

February 19 2020



A: Oblique coronal fat-suppressed T2-weighted MR image shows normal hypointense joint capsule at axillary recess (arrow). Note full-thickness tear of supraspinatus tendon (arrowheads) B: Oblique sagittal proton density MR image shows preserved subcoracoid fat triangle (asterisk) Credit: *American Journal of Roentgenology* (AJR)

Two MRI findings—joint capsule edema and thickness at the axillary recess, specifically—proved useful in predicting stiff shoulder in

patients with rotator cuff tears, according to an ahead-of-print article in the May issue of the *American Journal of Roentgenology (AJR)*.

Studying 106 [patients](#) with small to large (capsule edema and thickness in the axillary recess, Yoon Yi Kim of Korea's Veterans Health Service Medical Center assessed obliteration of the subcoracoid fat triangle, fatty degeneration of the torn rotator cuff muscle, and degree of retraction.

Tear size and location were determined by MRI findings and operative report, while associations between MRI findings and preoperative passive range of motion (ROM) were evaluated with simple and multiple linear regression analyses and proportional odds logistic regression analysis.

As Kim and colleagues wrote: "There was a significant, negative linear correlation between limited ROM at forward elevation and thickness of the joint capsule in the glenoid portion of the axillary recess ($p = 0.018$), external rotation and joint capsule edema in the humeral portion of the axillary recess ($p = 0.011$), and internal rotation and joint capsule [edema](#) in the glenoid portion of the axillary recess ($p = 0.007$)."

Fatty degeneration ($p = 0.003$) was an independent predictor of limited ROM on internal rotation. Meanwhile, male sex ($p = 0.041$) and posterosuperior rotator cuff tear ($p = 0.030$) were independent predictors of shoulder ROM on external rotation.

"This study is important," Kim et al. noted, "because it is the first to highlight joint capsule abnormality on MRI as a factor associated with stiff shoulder in patients with full-thickness rotator cuff tears."

More information: Yoon Yi Kim et al, MRI Findings Predictive of Shoulder Stiffness in Patients With Full-Thickness Rotator Cuff Tears,

American Journal of Roentgenology (2020). [DOI: 10.2214/AJR.19.21973](https://doi.org/10.2214/AJR.19.21973)

Provided by American Roentgen Ray Society

Citation: MRI findings predict shoulder stiffness for rotator cuff tears (2020, February 19)
retrieved 24 April 2024 from
<https://medicalxpress.com/news/2020-02-mri-shoulder-stiffness-rotator-cuff.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.