

Why obesity is more common than you think

February 20 2020, by John J Reilly



Credit: AI-generated image (disclaimer)

The World Health Organization has <u>described</u> obesity as a global epidemic and one of today's "most blatantly visible yet neglected public health problems." In the last few years <u>reports</u> on the rise of this lifethreatening condition among children and adults across the globe have been <u>alarming</u>.

A total of 107.7 million children and 603.7 million adults were



considered to be obese in 2015. But now our <u>research</u> shows that obesity is even more common than these reports suggest, and that in many countries most people are now living with obesity.

Obesity is a disease in which the amount of body fat is too high—high enough to harm health. It can increase the risk of diabetes, <u>high blood</u> <u>pressure</u>, <u>heart problems</u>, stroke and some cancers. But measuring body fatness accurately is difficult—and so most scientific studies and national surveys do not measure it, but use instead the weight-to-height <u>body mass index</u> (BMI) as a simple proxy for body fatness.

But the issue here is that the BMI can be a misleading proxy for high body fatness in both children and adults—and even more so in particular ethnic groups. This means that many people with apparently healthy BMI will actually be living with high body fatness without being aware of it, and so be at risk of the serious health consequences without realising it. Our research aimed to assess how problematic BMI is when used as a proxy for high body fatness in African children.

The problem with body-mass index

A high BMI or BMI-for-age—in children and adolescents BMI has to be adjusted for age because it increases with growth and maturation—is usually taken to represent high body fatness. The problem is that while almost all <u>adults</u> and <u>children</u> who have a high BMI will have a high level of body fat, many people with an apparently healthy BMI will also have a high level of body fat.

It is well known that BMI misclassifies the body fatness of athletes, bodybuilders and sportsmen and women, because a high muscle mass produces a high BMI. Only a small proportion of the population are athletic though, and the far more common problem with BMI is that it misclassifies body fatness of this much larger, non-athletic portion of the



population—but in the opposite direction, meaning it downplays its extent.

The extent of the misclassifying of body fatness by BMI matters. Most studies which have measured both body fatness and BMI have taken place in Europe and North America—and in these studies the prevalence of high body fatness has been as much as two to three times greater than the prevalence of high BMI. In one example 26% of US adults were obese according to their BMI, while 64% of the same adults were obese according to a more accurate measure which images the amount of fat in the body directly using X-rays.

Obesity and ethnicity

The problem of underestimating obesity is even worse in some population groups. The extent to which the BMI misclassifies the body fatness of people depends on their ethnicity. It is well established, but not well known, that BMI misclassifies children and adults in south and east Asia even more than Europeans.

There is emerging evidence that this problem extends to using BMI in other population groups too. In our <u>recent study</u> of children living in eight cities across Africa from Windhoek in the south to Tunis in the north, we found that 9% were obese according to their BMI-for-age, but 29% had high <u>body</u> fat according to a <u>stable isotope technique</u> designed for measuring <u>body fatness</u>.

By underestimating the prevalence of obesity because of reliance on BMI in clinics, research studies and national health surveys, we also underestimate the need for action to prevent and treat obesity. To use our African study as an example, the case for action at a 29% prevalence of obesity is much stronger than at a prevalence of 9%. Obesity is far more common than it seems, and it requires much greater effort and



more urgent attention than it is currently receiving.

As we enter a new decade it is time for national resolutions that are more ambitious. The main drivers of the global <u>obesity</u> epidemic are well known: excessive fat and sugar intake, lack of sleep, too much screen time, insufficient physical activity. We need global measures across all age groups to address these elements and we need them now.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Why obesity is more common than you think (2020, February 20) retrieved 4 May 2024 from https://medicalxpress.com/news/2020-02-obesity-common.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.