

Racial/ethnic disparities in stroke-prevention among patients undergoing dialysis

February 20 2020

In a study of patients with kidney failure and atrial fibrillation, racial/ethnic minorities experienced higher rates of stroke compared with non-Hispanic White patients, and they were less likely to fill prescriptions of stroke-preventive medications. The findings, which appear in an upcoming issue of *JASN*, indicate that equalizing the distribution of such prescriptions may help address stroke-related disparities among patients.

Because patients with kidney failure and [atrial fibrillation](#), or an [irregular heartbeat](#), are at risk of developing strokes, they may benefit from taking blood thinners as a preventive measure. Prior studies of patients with atrial fibrillation, with or without kidney failure, have reported that minority racial/[ethnic groups](#) face a higher risk of stroke compared with Whites.

To investigate the potential causes of such disparities, a team led by Paul L. Kimmel, MD (National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health) examined information from a national registry of patients with kidney failure.

For the study, the researchers analyzed information from the United States Renal Data System to identify patients with [kidney failure](#) who initiated hemodialysis from 2006 to 2013, and then they identified those with a subsequent atrial fibrillation diagnosis and Medicare Part A/B/D insurance coverage.

Among 56,587 patients who were followed for one year, the number of strokes per 1,000 people were 84, 94, 97, and 102 in non-Hispanic White, Black, Hispanic White, and Asian patients, respectively.

Black, Hispanic White, and Asian patients filled prescriptions of the blood-thinner warfarin less often than non-Hispanic White patients, and they were more likely to experience stroke. Specifically, Black, Hispanic White, and Asian patients were 10%, 17%, and 28% less likely than non-Hispanic Whites to fill a warfarin prescription, respectively, and they were 13%, 15%, and 16% more likely to experience stroke. (The database did not provide information on whether patients were prescribed drugs, only whether they filled prescriptions.)

Analyses suggested that equalizing the warfarin distribution to that in the non-Hispanic White patient population would prevent 7%, 10%, and 12% of the stroke disparity among Black, Hispanic White, and Asian patients, respectively.

"Institutional features and medical choices may play key roles underlying differences between prescriptions received by members of different groups. Our estimates suggest equalization of prescription of warfarin, a relatively inexpensive anticoagulant medication, across all ethnic/[racial groups](#) would be associated with decreased stroke rates in Black, Asian, and Hispanic White patients on hemodialysis," said Dr. Kimmel.

More information: "Racial/Ethnic Disparities in Atrial Fibrillation Treatment and Outcomes among Dialysis Patients in the United States," [DOI: 10.1681/ASN.2019050543](https://doi.org/10.1681/ASN.2019050543)

Provided by American Society of Nephrology

Citation: Racial/ethnic disparities in stroke-prevention among patients undergoing dialysis (2020, February 20) retrieved 26 April 2024 from <https://medicalxpress.com/news/2020-02-raciaethnic-disparities-stroke-prevention-patients-dialysis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.