

Developing seizures after stroke may increase risk of death, disability

February 12 2020

Seizures may be linked to a higher risk of death or disability in adults who have had a severe ischemic stroke, according to preliminary research to be presented at the American Stroke Association's International Stroke Conference 2020—Feb. 19-21 in Los Angeles, a world premier meeting for researchers and clinicians dedicated to the science of stroke and brain health.

Ischemic stroke occurs when blood vessels leading to the brain are blocked. Seizures are abnormal brain activity that may cause uncontrollable shaking movements, loss of consciousness and confusion. Ischemic [stroke patients](#) who had seizures within seven days after their stroke were at higher risk of [seizure](#) recurrence and of being diagnosed with epilepsy.

"In the United States, stroke is the most common cause of epilepsy in adults. In Mexico, stroke is now the leading cause of epilepsy in adults as well," said Erwin Chiquete, M.D., Ph.D., lead study author and a neurologist and researcher at The Salvador Zubiran National Institute of Medical Sciences and Nutrition in Mexico City. "However, we suspect that seizures and epilepsy in stroke patients are still under-recognized by physicians."

In this study, researchers identified [risk factors](#) that predict the possibility of developing seizures or epilepsy (recurring seizures) following a [severe stroke](#), and then assessed the impact of post-stroke seizures on death or disability up to one year after stroke.

Researchers analyzed electronic medical records data of 1,246 [hospital patients](#) without epilepsy upon admission, aged 18 to 94 (85% over age 40) and treated for [acute ischemic stroke](#) in 59 centers located in Mexico. After following the patients for 12 months, researchers found:

- 8% of stroke survivors suffered acute seizures within a week of their stroke, and almost 5% of these patients had at least one more seizure during the 12-month follow-up period.
- Patients 18 to 40 years old were the most vulnerable group of stroke survivors. More than 13% of these patients developed recurrent seizures and were diagnosed with epilepsy.
- Patients younger than 65 were 69% more likely to develop seizures than older patients.
- The larger the injury to the brain from the stroke, the greater the likelihood of having a seizure. The odds of developing seizures were more than two times higher in people with severe brain injury from stroke and in those who had a second stroke.
- Those who developed acute seizures were 37% more likely to die or develop physical and/or mental disabilities within one year following their stroke.
- Other factors that significantly increased risk of death or disability included being over age 65, severity of stroke, heart failure and atrial fibrillation.

"We think that the secondary prevention of ischemic stroke and controlling seizures with anti-epileptic medication during the twelve months after stroke are very important to prevent death and to decrease disability," Chiquete said. "Our research suggests that severe [ischemic stroke](#) can cause epilepsy in a higher percentage of patients than we originally thought or has been shown in previous studies."

Provided by American Heart Association

Citation: Developing seizures after stroke may increase risk of death, disability (2020, February 12) retrieved 25 April 2024 from <https://medicalxpress.com/news/2020-02-seizures-death-disability.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.