


For weight-loss surgery patients who quit smoking, relapse is common

February 21 2020



Source: King, et al., *Annals of Surgery*, 2020.

Most adult smokers who undergo weight-loss surgery quit in the month before their operation. But the majority relapse and, within seven years, smoking rates climb to pre-surgery levels. Credit: UPMC

Although 1 in 7 adults smoke cigarettes the year prior to undergoing weight-loss surgery, nearly all successfully quit at least a month before their operation. However, smoking prevalence steadily climbs to pre-surgery levels within seven years, according to new research led by the University of Pittsburgh Graduate School of Public Health.

The findings—reported today in the *Annals of Surgery*—suggest that there may be missed opportunities to engage patients in interventions to improve long-term smoking cessation rates, particularly at regular post-[surgery](#) checkups.

"Smoking cessation prior to surgery is strongly recommended to reduce surgical complications," said lead author Wendy King, Ph.D., associate professor of epidemiology at Pitt Public Health. "But there isn't the same emphasis on maintaining cessation after surgery. Our findings show that there is a need for ongoing support in order to reduce and quickly respond to relapses."

King and her colleagues followed 1,770 adults who underwent Roux-en-Y [gastric bypass surgery](#)—a procedure that reduces the size of the stomach and bypasses part of the small intestine—for seven years post-surgery, annually surveying them about their smoking habits. The participants were enrolled in the National Institutes of Health-funded Longitudinal Assessment of Bariatric Surgery-2 (LABS-2), a prospective, observational study of patients undergoing [weight-loss surgery](#) at one of 10 hospitals across the United States.

More than 45% of the participants reported a history of smoking prior to surgery, with 14% still smoking in the year before surgery, which fell to 2% in the month before surgery. The rate rebounded to nearly 10% in the year following surgery and steadily climbed back to 14% by seven years post-surgery.

"Interestingly, the people who picked up smoking post-surgery weren't just the people who quit smoking in the year prior to surgery, presumably to prepare for the operation. Many had never smoked to begin with," said co-author Gretchen White, Ph.D., assistant professor in Pitt's School of Medicine, explaining that 2 out of 5 people who smoked after surgery had quit more than a year before their operation or hadn't ever smoked.

Additionally, people who identified as smokers post-surgery smoked more, going from an average of a dozen cigarettes per day in the year before surgery to more than 15 cigarettes per day seven years post-surgery. These findings contrast with concurrent reductions in smoking prevalence and intensity in the general U.S. population.

The researchers hypothesized that weight control would be a key reason weight-loss patients took up smoking after surgery, but found that the prevalence of smoking for weight control was actually fairly stable over time, at about 2% pre- and post-surgery, and did not appear to be related to smoking more cigarettes. King noted that "this surprised everyone, as there is a general assumption that weight control is a main motivator for smoking."

While the study was not designed to find a biological reason for the results, the researchers did observe that gastric bypass patients were more likely to smoke post-surgery than patients who underwent gastric banding, where a silicone belt is surgically inserted around the stomach to reduce the amount of food it can hold. A recent study showed that gastric bypass increases exposure to the psychoactive nicotine metabolite cotinine. Just as gastric bypass increases the risk of alcohol use disorder due to changes in alcohol metabolism that lead to higher and quicker elevation of blood alcohol levels, it may also increase risk of smoking via nicotine metabolism, King suggested.

The scientists identified several factors that predict which patients would be most likely to take up smoking after surgery. Not surprisingly, a prior history of [smoking](#) was the greatest risk factor. In addition, younger age, poverty, being married or living as married, and drug use were each associated with increased risk.

More information: Wendy C. King et al, Changes in Smoking Behavior Before and After Gastric Bypass, *Annals of Surgery* (2020).
[DOI: 10.1097/SLA.0000000000003828](https://doi.org/10.1097/SLA.0000000000003828)

Provided by University of Pittsburgh

Citation: For weight-loss surgery patients who quit smoking, relapse is common (2020, February 21) retrieved 17 May 2024 from
<https://medicalxpress.com/news/2020-02-weight-loss-surgery-patients-relapse-common.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--